Journal of Functional Analysis 269 (2015) 1714–1768

Contents lists available at ScienceDirect

Journal of Functional Analysis

www.elsevier.com/locate/jfa

Long time behavior of solutions of Fisher-KPP equation with advection and free boundaries ☆

Hong Gu [a], Bendong Lou [b,*], Maolin Zhou [c]

[a] *School of Applied Mathematics, Nanjing University of Finance & Economics, Nanjing 210023, China*
[b] *Department of Mathematics, Tongji University, Shanghai 200092, China*
[c] *School of Science and Technology, University of New England, Armidale, NSW 2351, Australia*

ARTICLE INFO

Article history:
Received 8 October 2014
Accepted 5 July 2015
Available online 21 July 2015
Communicated by F.-H. Lin

MSC:
35K20
35K55
35R35
35B40

Keywords:
Fisher-KPP equation
Advection
Free boundary problem
Long time behavior

ABSTRACT

We consider Fisher-KPP equation with advection: $u_t = u_{xx} - \beta u_x + f(u)$ for $x \in (g(t), h(t))$, where $g(t)$ and $h(t)$ are two free boundaries satisfying Stefan conditions. This equation is used to describe the population dynamics in advective environments. We study the influence of the advection coefficient $-\beta$ on the long time behavior of the solutions. We find two parameters c_0 and β^* with $\beta^* > c_0 > 0$ which play key roles in the dynamics, here c_0 is the minimal speed of the traveling waves of Fisher-KPP equation. More precisely, by studying a family of the initial data $\{\sigma\phi\}_{\sigma>0}$ (where ϕ is some compactly supported positive function), we show that: (1) in case $\beta \in (0, c_0)$, there exists $\sigma^* \geqslant 0$ such that spreading happens when $\sigma > \sigma^*$ (i.e., $u(t, \cdot\,; \sigma\phi) \to 1$ locally uniformly in $\mathbb{R}$) and vanishing happens when $\sigma \in (0, \sigma^*]$ (i.e., $[g(t), h(t)]$ remains bounded and $u(t, \cdot\,; \sigma\phi) \to 0$ uniformly in $[g(t), h(t)]$); (2) in case $\beta \in (c_0, \beta^*)$, there exists $\sigma^* > 0$ such that virtual spreading happens when $\sigma > \sigma^*$ (i.e., $u(t, \cdot\,; \sigma\phi) \to 0$ locally uniformly in $[g(t), \infty)$ and $u(t, \cdot + ct; \sigma\phi) \to 1$ locally uniformly in $\mathbb{R}$ for some $c > \beta - c_0$), vanishing happens when $\sigma \in (0, \sigma^*)$, and in the transition case $\sigma = \sigma^*$, $u(t, \cdot + o(t); \sigma\phi) \to V^*(\cdot - (\beta - c_0)t)$ uniformly, the latter is a traveling wave with a "big head" near the free

☆ This research was partly supported by NSFC (No. 11271285).
* Corresponding author.
E-mail addresses: honggu87@126.com (H. Gu), blou@tongji.edu.cn (B. Lou), mzhou6@une.edu.au (M. Zhou).

http://dx.doi.org/10.1016/j.jfa.2015.07.002
0022-1236/© 2015 Elsevier Inc. All rights reserved.

boundary $x = (\beta - c_0)t$ and with an infinite long "tail" on the left; (3) in case $\beta = c_0$, there exists $\sigma^* > 0$ such that virtual spreading happens when $\sigma > \sigma^*$ and $u(t, \cdot; \sigma\phi) \to 0$ uniformly in $[g(t), h(t)]$ when $\sigma \in (0, \sigma^*]$; (4) in case $\beta \geqslant \beta^*$, vanishing happens for any solution.
© 2015 Elsevier Inc. All rights reserved.

1. Introduction

In this paper, we consider the following problem

$$\begin{cases} u_t = u_{xx} - \beta u_x + f(u), & g(t) < x < h(t),\ t > 0, \\ u(t, g(t)) = 0, \quad g'(t) = -\mu u_x(t, g(t)), & t > 0, \\ u(t, h(t)) = 0, \quad h'(t) = -\mu u_x(t, h(t)), & t > 0, \\ -g(0) = h(0) = h_0, \quad u(0, x) = u_0(x), & -h_0 \leqslant x \leqslant h_0, \end{cases} \tag{P}$$

where μ and β are positive constants, $h_0 > 0$ and u_0 is the initial data to be specified in the next section, $f : [0, \infty) \to \mathbb{R}$ is a C^1 function satisfying

$$\begin{cases} f(0) = f(1) = 0, \quad (1 - u)f(u) > 0 \text{ for } u > 0 \text{ and } u \neq 1, \\ f'(0) > 0,\ f'(1) < 0 \text{ and } f(u) \leqslant f'(0)u \text{ for } u \geqslant 0. \end{cases} \tag{F}$$

The problem (P) is used to model the spreading of a new or invasive species, under the influence of diffusion and advection. The unknown $u(t, x)$ denotes the population density over a one-dimensional habitat and the free boundaries $x = g(t)$ and $x = h(t)$ represent the expanding fronts of the species. We assume that the free boundaries move according to one-phase Stefan condition, which is a kind of free boundary conditions widely used in the study of melting of ice [34], wound healing [9], and population dynamics [7,12,13]. The derivation of one-phase or two-phase Stefan conditions in population models as singular limits of competition–diffusion systems can be found in [22,23] etc.

When $\beta = 0$ (i.e., there is no advection in the environment), the qualitative properties of the problem (P) were studied by Du and Lin [12] for logistic nonlinearity $f(u) = u(1 - u)$. Among others, they proved that, when $h_0 \geqslant \frac{\pi}{2}$, any solution of (P) with $\beta = 0$ grows up and converges to 1 (which is called *spreading* phenomenon); when $h_0 < \frac{\pi}{2}$, spreading happens if μ is large and *vanishing* happens if μ is small (i.e., the solution converges to 0). The vanishing phenomenon is a remarkable result since it shows that the presence of free boundaries may avoid the so-called *hair-trigger effect*, which is a phenomenon shown in [2]: spreading always happens for a solution of the Cauchy problem for $u_t = u_{xx} + f(u)$, no matter how small the positive initial data is. Recently, Du and Lou [13] extended the results in [12] to the problem with general monostable, bistable and combustion types of f, and gave a rather complete description on the long time behavior of the solutions. In addition, Kaneko and Yamada [27], Liu and Lou [28,29] studied the problem (P) with $\beta = 0$ and with a fixed boundary $g(t) \equiv 0$, Du and Guo

[11,10], Du, Matano and Wang [15], Zhou and Xiao [39], Wang [38] studied the problem (without advection) in higher dimension spaces and/or in spatial heterogeneous environments. Besides the qualitative properties, another interesting problem is the asymptotic spreading speeds of the free boundaries when spreading happens. Du and Lin [12], Du and Lou [13] proved that, when spreading happens for a solution (u, g, h) of the problem (P) with $\beta = 0$,

$$c^* := \lim_{t\to\infty} \frac{h(t)}{t} = \lim_{t\to\infty} \frac{-g(t)}{t} > 0. \tag{1}$$

Recently, Du, Matsuzawa and Zhou [16] improved this result to better ones:

$$\lim_{t\to\infty} h'(t) = \lim_{t\to\infty} [-g'(t)] = c^*, \quad \lim_{t\to\infty} [h(t) - c^*t] = H_\infty, \quad \lim_{t\to\infty} [g(t) + c^*t] = G_\infty, \tag{2}$$

for some $H_\infty, G_\infty \in \mathbb{R}$.

In this paper we consider the problem (P) with $\beta > 0$, which means that the spreading of a species is affected by advection. In the field of ecology, organisms can often sense and respond to local environmental cues by moving towards favorable habitats, and these movements usually depend upon a combination of local biotic and abiotic factors such as stream, climate, food and predators. For example, some diseases spread along the wind direction. In 2009, Maidana and Yang [30] studied the propagation of West Nile Virus from New York City to California state. It was observed that West Nile Virus appeared for the first time in New York City in the summer of 1999. In the second year the wave front travels 187 km to the north and 1100 km to the south. Therefore, they took account of the advection movement and showed that bird advection becomes an important factor for lower mosquito biting rates. Another example is that Averill [3] considered the effect of intermediate advection on the dynamics of two-species competition system, and provided a concrete range of advection strength for the coexistence of two competing species. Moreover, three different kinds of transitions from small advection to large advection were illustrated theoretically and numerically. Many other examples involving advection can also be found in the field of ecology (cf. [4,5,8,32,33,35–37] etc.).

From a mathematical point of view, to involve the influence of advection, one of the simplest but probably still realistic approaches is to assume that species can move up along the gradient of the density, as considered in [6,24,25,33,35–37] etc.

Gu, Lin and Lou [19,20] studied the problem (P) with small advection. They proved a spreading–vanishing dichotomy result on the long time behavior of positive solutions of (P), which is similar to the conclusions in [12,13] for equations without advection. They also proved that, when spreading happens for a solution of (P) with small advection, its rightward spreading speed is bigger than the leftward one:

$$\lim_{t\to\infty} \frac{h(t)}{t} > \lim_{t\to\infty} \frac{-g(t)}{t} > 0. \tag{3}$$

Recently, Kaneko and Matsuzawa [26] improved this result to some conclusions like (2).

Our main purpose in this paper is to study the influence of the advection term $-\beta u_x$ on the long time behavior of solutions of (P). As we will see below, our study improves the results in [19,20,26] since we will study the problem (P) for all $\beta > 0$, not only for small β. Especially, when β is large, the phenomenon is much more complicated and more interesting than the case where β is small.

To sketch the influence of β, we introduce two important traveling waves. First, consider the following problem

$$\begin{cases} q''(z) - cq'(z) + f(q) = 0, \quad z \in \mathbb{R}, \\ q(-\infty) = 0,\ q(+\infty) = 1,\ q(0) = 1/2, \quad q'(z) > 0 \text{ for } z \in \mathbb{R}. \end{cases} \tag{4}$$

It is well known that this problem has a solution $q(z;c)$ if and only if $c \geqslant c_0$, where

$$c_0 := 2\sqrt{f'(0)}$$

is called the minimal speed of the traveling waves of Fisher-KPP equation. Denote $Q(z) := q(z;c_0)$, then $u(t,x) = Q(x - (\beta - c_0)t)$ is a traveling wave of $u_t = u_{xx} - \beta u_x + f(u)$. It travels leftward (resp. rightward) if and only if $\beta < c_0$ (resp. $\beta > c_0$). Next we consider the following problem

$$\begin{cases} q''(z) + (c-\beta)q'(z) + f(q) = 0, \quad z \in (-\infty, 0), \\ q(0) = 0,\ q(-\infty) = 1,\ -\mu q'(0) = c, \quad q'(z) < 0 \text{ for } z \in (-\infty, 0]. \end{cases} \tag{5}$$

As is shown in Lemma 3.4 (see also [12,13,20]), for any $\beta > 0$, this problem has a unique solution $(c,q) = (c^*, U^*(z))$. So $u(t,x) = U^*(x - c^*t)$ is a solution of $u_t = u_{xx} - \beta u_x + f(u)$, with $u(t, c^*t) = 0$, $c^* = -\mu u_x(t, c^*t)$. It is called a *traveling semi-wave* in [13] since it is only defined for $x \leqslant c^*t$. We also write c^* as $c^*(\beta)$ to emphasize the dependence of c^* on β, then we will show in Lemma 3.4 that the equation $\beta - c_0 = c^*(\beta)$ has a unique root $\beta^* > c_0$:

$$\beta^* - c_0 = c^*(\beta^*). \tag{6}$$

We will see below that the traveling wave $Q(x - (\beta - c_0)t)$ and the traveling semi-wave $U^*(x - c^*t)$ are of special importance in the study of spreading solutions. To explain their roles intuitively, we consider the problem (P) with initial data $u_0(x)$ which is even and

$$u_0(x) = \begin{cases} 1, & x \in [0, h_0 - 1], \\ \text{smooth and decreasing}, & x \in [h_0 - 1, h_0], \end{cases} \qquad \text{with } h_0 \gg 1.$$

It is easily seen by the maximum principle that $u(t,\cdot)$ has exactly one maximum point. As usual, we call the sharp decreasing part in the graph of $u(t,\cdot)$ the *front*, and call the sharp increasing part on the left side the *back*. Now we sketch the influence of the advection $-\beta u_x$. *Case 1:* When $\beta \in (0, c_0)$, the advection influence is not strong, the solution has enough space between the back and the front to grow up and to converge

to 1. Its front approaches a profile like $U^*(\cdot)$ and moves rightward at a speed $\approx c^*$. Its back approaches a profile like $U^*(-\cdot)$ and moves leftward at a speed smaller than $c_0-\beta$ (see details in Theorem 2.5 below). This case is similar as the spreading phenomena in [12,13] for the equation with $\beta=0$. *Case 2:* When $\beta\in(c_0,\beta^*)$ with β^* being the unique root of (6), the traveling wave $Q(x-(\beta-c_0)t)$ travels rightward at a speed $\beta-c_0>0$. Hence the back of the solution u, with a shape like $Q(x-(\beta-c_0)t)$, is pushed by $Q(x-(\beta-c_0)t)$ to move rightward at a speed $\approx\beta-c_0$, and so $u\to 0$ locally uniformly. But, when the initial domain is wide enough, the solution still has enough space to grow up between the back and the front since the front moves rightward (at a speed $\approx c^*$) faster than the back. In this paper we call such a phenomenon as *virtual spreading* (see Theorem 2.2 and Lemma 4.11 below). *Case 3:* When $\beta=c_0$, the traveling wave $Q(x-(\beta-c_0)t)=Q(x)$ is indeed a stationary solution of $(P)_1$ (hereafter, $(P)_1$ denotes the equation $u_t=u_{xx}-\beta u_x+f(u)$ in (P)). However, the back of the solution u still moves rightward at a speed $O(t^{-1})$ since it starts from a compactly supported initial data u_0 (cf. [21] and see details below). Hence virtual spreading still happens when h_0 is sufficiently large, since the front moves rightward at speed c^*, faster than the back. *Case 4:* When $\beta>\beta^*$, the back moves rightward (at a speed $\approx\beta-c_0$) faster than the front (which moves rightward at speed $\approx c^*<\beta-c_0$). So the solution is suppressed by its back, and then $u\to 0$ uniformly. In summary, the long time behavior of the solutions is quite different for $\beta\in(0,c_0)$, $\beta=c_0$, $\beta\in(c_0,\beta^*)$ and $\beta>\beta^*$.

This paper is organized as follows. In Section 2 we present our main results. In Section 3 we give some preliminaries including the comparison principles, stationary solutions, several types of traveling waves, zero number arguments and some upper bound estimates. In Section 4 we study the influence of the advection on the long time behavior of the solutions. In Section 5, we revisit the virtual spreading phenomena and give a uniform convergence for such solutions.

2. Main results

Throughout this paper we choose initial data u_0 from the following set:

$$\mathscr{X}(h_0):=\big\{\phi\in C^2([-h_0,h_0]):\phi(-h_0)=\phi(h_0)=0,\ \phi'(-h_0)>0,\\ \phi'(h_0)<0,\ \phi(x)>0 \text{ in } (-h_0,h_0)\big\},\tag{7}$$

where $h_0>0$ is any given real number. By a similar argument as in [12,13], one can show that, for any initial data $u_0\in\mathscr{X}(h_0)$, the problem (P) has a time-global solution (u,g,h), with $u\in C^{1+\nu/2,2+\nu}(\Omega)$ and $g,h\in C^{1+\nu/2}((0,\infty))$ for any $\nu\in(0,1)$, where $\Omega:=\{(t,x):x\in[g(t),h(t)],\ t\in(0,\infty)\}$. Moreover, it follows from the maximum principle that, when $t>0$, the solution u is positive in $(g(t),h(t))$, $u_x(t,g(t))>0$ and $u_x(t,h(t))<0$. Hence $g'(t)<0$, $h'(t)>0$. Denote

$$g_\infty:=\lim_{t\to\infty}g(t),\quad h_\infty:=\lim_{t\to\infty}h(t),\quad I(t):=[g(t),h(t)]\quad\text{and}\quad I_\infty:=(g_\infty,h_\infty).$$

In what follows, we mainly consider the solution of (P) with initial data $u_0 = \sigma\phi$ for some given $\phi \in \mathscr{X}(h_0)$ and $\sigma \geqslant 0$. We also use $(u(t,x;\sigma\phi), g(t;\sigma\phi), h(t;\sigma\phi))$ to denote such a solution. Now we list some possible situations for the solutions of (P):

- *spreading*: $I_\infty = \mathbb{R}$ and

$$\lim_{t\to\infty} u(t,\cdot) = 1 \text{ locally uniformly in } \mathbb{R}; \tag{8}$$

- *vanishing*: I_∞ is a bounded interval and

$$\lim_{t\to\infty} \max_{g(t)\leqslant x\leqslant h(t)} u(t,x) = 0; \tag{9}$$

- *virtual spreading*: $g_\infty > -\infty$, $h_\infty = +\infty$,

$$\lim_{t\to\infty} u(t,\cdot) = 0 \text{ locally uniformly in } I_\infty \tag{10}$$

  and

$$\lim_{t\to\infty} u(t,\cdot + ct) = 1 \text{ locally uniformly in } \mathbb{R}, \quad \text{for some } c > 0; \tag{11}$$

- *virtual vanishing*: $g_\infty > -\infty$, $h_\infty = +\infty$ and (9) holds.

When the advection is small, we have the following conclusion on the long time behavior of the solutions.

Theorem 2.1 *(The case $\beta \in (0, c_0)$). Assume $0 < \beta < c_0$ and (u, g, h) is a time-global solution of (P) with initial data $u_0 = \sigma\phi$ for some $\phi \in \mathscr{X}(h_0)$. Then there exists $\sigma^* = \sigma^*(h_0, \phi) \in [0, \infty]$ such that*

(i) vanishing *happens when $\sigma \in [0, \sigma^*]$, with $|I_\infty| = h_\infty - g_\infty \leqslant \frac{2\pi}{\sqrt{c_0^2-\beta^2}}$;*
(ii) spreading *happens when $\sigma > \sigma^*$.*

From this theorem we see that the long time behavior of the solutions of (P) with small advection: $\beta \in (0, c_0)$ is similar as the case without advection: $\beta = 0$ (cf. [12,13,19]). The main reason is that in both cases the problem (P) has exactly two stationary solutions: 0 and 1 in $\mathbb{R}$. The proof of this theorem, which is given in Section 4.2, is also similar as that for $\beta = 0$.

Next we consider the case where the advection is not small: $\beta \geqslant c_0$. The most interesting phenomenon appears in the problem with medium-sized advection: $\beta \in [c_0, \beta^*)$, where β^* is the unique root of (6).

Theorem 2.2 *(The case $c_0 < \beta < \beta^*$). Assume $c_0 < \beta < \beta^*$ and (u, g, h) is a time-global solution of (P) with initial data $u_0 = \sigma\phi$ for some $\phi \in \mathscr{X}(h_0)$. Then there exists $\sigma^* = \sigma^*(h_0, \phi) \in (0, \infty]$ such that*

(i) virtual spreading *happens when $\sigma > \sigma^*$, and*

$$\lim_{t\to\infty} u(t, \cdot + ct) = 1 \text{ locally uniformly in } \mathbb{R}, \quad \text{for any } c \in (\beta - c_0, c^*),$$

where $c^* = c^*(\beta)$ is the speed of the traveling semi-wave in (5);

(ii) vanishing *happens when $0 < \sigma < \sigma^*$;*

(iii) *in the* transition *case $\sigma = \sigma^*$: $g_\infty > -\infty$, $h_\infty = +\infty$,*

$$\lim_{t\to\infty} h'(t) = \beta - c_0 \quad \text{and} \quad h(t) = (\beta - c_0)t + \varrho(t)$$

with $\varrho(t) = o(t)$ and $\varrho(t) \to \infty$ $(t \to \infty)$. In addition,

$$\lim_{t\to\infty} \|u(t, \cdot) - V^*(\cdot - (\beta - c_0)t - \varrho(t))\|_{L^\infty(I(t))} = 0, \tag{12}$$

where $V^*(z)$ is the unique solution of

$$\begin{cases} q''(z) - c_0 q'(z) + f(q) = 0 & \text{for } z \in (-\infty, 0), \\ q(0) = 0,\ q(-\infty) = 0,\ q(z) > 0 \text{ for } z \in (-\infty, 0),\ -\mu q'(0) = \beta - c_0. \end{cases} \tag{13}$$

In the next section we will see that V^* has a *tadpole-like* shape: it has a "big head" and a boundary on the right side and an infinite long "tail" on the left side. So we call $V^*(x - (\beta - c_0)t)$ a *tadpole-like traveling wave* with speed $\beta - c_0$, which exists if and only if $\beta \in (c_0, \beta^*)$ (see Lemma 3.5 below). Theorem 2.2(iii) implies that, roughly, $u(t, x)$ converges to this traveling wave.

In Aronson and Weinberger [2], it was shown that any positive solution of the Cauchy problem for Fisher-KPP equation converges to 1 (i.e., *hair-trigger effect*). In [12,13], by introducing the free boundaries, the authors proved a spreading–vanishing dichotomy on the long time behavior of the solutions of Fisher-KPP equation. In particular, vanishing may happen for some solutions. Now our Theorem 2.2 gives the third possibility besides the virtual spreading and vanishing, that is, with a medium-sized advection in the equation, there may exist a transition state: the solution converges to a *tadpole-like traveling wave*. This interesting phenomenon is new comparing with the results for Cauchy problems and for free boundary problems without advection.

Theorem 2.3 *(The case $\beta = c_0$). Assume $\beta = c_0$ and (u, g, h) is a time-global solution of (P) with initial data $u_0 = \sigma\phi$ for some $\phi \in \mathscr{X}(h_0)$. Then there exist $\sigma_*, \sigma^* \in (0, \infty]$ with $\sigma_* \leqslant \sigma^*$ such that*

(i) virtual spreading *happens when* $\sigma > \sigma^*$, *and*

$$\lim_{t\to\infty} u(t, \cdot + ct) = 1 \text{ locally uniformly in } \mathbb{R}, \quad \text{for any } c \in (0, c^*),$$

where $c^* = c^*(\beta)$ *is the speed of the traveling semi-wave in* (5);

(ii) vanishing *happens when* $0 < \sigma < \sigma_*$;

(iii) virtual vanishing *happens when* $\sigma \in [\sigma_*, \sigma^*]$.

The transition cases in Theorem 2.2 and Theorem 2.3 are different. In case $c_0 < \beta < \beta^*$, a solution $u(t, x; \sigma\phi)$ is a transition one only if the initial value takes the sharp threshold value $\sigma^*\phi$. However, in case $\beta = c_0$, we obtain transition solutions whose initial data are taken from $\{\sigma\phi \mid \sigma \in [\sigma_*, \sigma^*]\}$. Whether or not this domain is a singleton: $\sigma_* = \sigma^*$ is still open now. The difficulty in studying this problem is that virtual vanishing solutions have no "shapes", so it is not easy to compare one to another.

The conclusions for the problem with large advection: $\beta \geqslant \beta^*$ is rather simple.

Theorem 2.4 *(The case* $\beta \geqslant \beta^*$*). Assume* $\beta \geqslant \beta^*$ *and* (u, g, h) *is a time-global solution of* (P) *with initial data* $u_0 \in \mathscr{X}(h_0)$*. Then vanishing happens.*

Besides the convergence/dichotomy/trichotomy results on the long time behavior of the solutions as stated in the previous theorems, we can say more about the solutions when (virtual) spreading happens. It turns out that, when $\beta \in [c_0, \beta^*)$, the virtual spreading solution can be characterized by the rightward traveling semi-wave $U^*(x - c^*t)$ and the traveling wave $Q(x - (\beta - c_0)t)$; when $\beta \in (0, c_0)$, the spreading solution can be characterized by $U^*(x - c^*t)$ and the leftward traveling semi-wave $U_l^*(x - c_l^*t)$. Here (c_l^*, U_l^*) (with $c_l^* < 0$) is the unique solution of the following problem with $\beta \in (0, c_0)$ (see details in Section 3.3)

$$\begin{cases} q''(z) + (c - \beta)q'(z) + f(q) = 0, & z \in (0, \infty), \\ q(0) = 0,\ q(\infty) = 1,\ -\mu q'(0) = c,\ q'(z) > 0 \text{ for } z \in (0, \infty). \end{cases} \tag{14}$$

Using these traveling waves we can give the asymptotic profiles for (virtual) spreading solutions.

Theorem 2.5. *Assume spreading or virtual spreading happens for a solution of* (P) *as in Theorems 2.1, 2.2 or 2.3. Let* (c^*, U^*) *be the unique solution of* (5) *with* $c^* > 0$.

(i) When $\beta \in (0, c_0)$*, let* (c_l^*, U_l^*) *be the unique solution of* (14) *with* $0 < -c_l^* < c^*$*. Then there exist* $H_\infty, G_\infty \in \mathbb{R}$ *such that*

$$\lim_{t\to\infty} [h(t) - c^*t] = H_\infty, \quad \lim_{t\to\infty} h'(t) = c^*, \tag{15}$$

$$\lim_{t\to\infty} [g(t) - c_l^*t] = G_\infty, \quad \lim_{t\to\infty} g'(t) = c_l^*, \tag{16}$$

and, if we extend U^*, U_l^* to be zero outside their supports we have

$$\lim_{t\to\infty} \left\| u(t,\cdot) - U^*(\cdot - c^*t - H_\infty)\cdot U_l^*\big(\cdot - c_l^*t - G_\infty\big)\right\|_{L^\infty(I(t))} = 0. \tag{17}$$

(ii) When $\beta \in [c_0, \beta^*)$, (15) holds for some $H_\infty \in \mathbb{R}$. Moreover, if we extend U^* to be zero outside its support, then

$$\lim_{t\to\infty} \left\| u(t,\cdot) - U^*(\cdot - c^*t - H_\infty)\cdot Q\big(\cdot - (\beta - c_0)t - \theta(t)\big)\right\|_{L^\infty(I(t))} = 0 \tag{18}$$

for some function $\theta(t)$ satisfying $\theta(t) = o(t)$ and $\theta(t) \to \infty$ $(t \to \infty)$.

Assume $\beta \in [0, c_0)$ and spreading happens for a solution (u, g, h) of (P). The asymptotic spreading speed $\lim_{t\to\infty} \frac{h(t)}{t} = c^*$ was obtained in [12,13] for the case $\beta = 0$, and in [20] for the case $\beta \in (0, c_0)$. Recently, Du, Matsuzawa and Zhou [16], Kaneko and Matsuzawa [26] improved them to analogues of (15), (16) and (17). Note that our theorem includes both the case $\beta \in (0, c_0)$ and the case $\beta \in [c_0, \beta^*)$. The proof of (18) will be given in the last section, based on the fact that Q is steeper than any other entire solution (see Section 5 below and [17]).

3. Preliminaries

In this section we first give some comparison principles and then present all the bounded stationary solutions and traveling wave solutions of $(P)_1$ which will be used for comparison. In the fourth subsection we give some results on the zero numbers of the solutions of linear equations which will play key roles in our approach. In the last subsection we give some precise upper bound estimates for the solutions.

3.1. The comparison principle

In this subsection we give two types of comparison principles which will be used frequently in this paper. Similar as in [12,13], we have

Lemma 3.1. *Assume $T \in (0, \infty)$, $\bar{g}(t), \bar{h}(t) \in C^1([0, T])$, $\bar{u}(t, x) \in C(\overline{D}_T) \cap C^{1,2}(D_T)$ with $D_T = \{(t, x) \in \mathbb{R}^2 \mid 0 < t \leqslant T,\ \bar{g}(t) < x < \bar{h}(t)\}$, and*

$$\begin{cases} \bar{u}_t \geqslant \bar{u}_{xx} - \beta\bar{u}_x + f(\bar{u}), & 0 < t \leqslant T,\ \bar{g}(t) < x < \bar{h}(t), \\ \bar{u} = 0, \quad \bar{g}'(t) \leqslant -\mu\bar{u}_x, & 0 < t \leqslant T,\ x = \bar{g}(t), \\ \bar{u} = 0, \quad \bar{h}'(t) \geqslant -\mu\bar{u}_x, & 0 < t \leqslant T,\ x = \bar{h}(t). \end{cases}$$

If

$$[-h_0, h_0] \subset [\bar{g}(0), \bar{h}(0)] \quad \textit{and} \quad u_0(x) \leqslant \bar{u}(0, x) \textit{ for } x \in [-h_0, h_0],$$

and (u, g, h) *is a solution of* (P)*, then*

$$g(t) \geqslant \bar{g}(t),\ h(t) \leqslant \bar{h}(t) \quad \text{for } t \in (0, T],$$
$$u(t, x) \leqslant \bar{u}(t, x) \quad \text{for } t \in (0, T] \text{ and } x \in (g(t), h(t)).$$

Lemma 3.2. *Assume* $T \in (0, \infty)$, $l(t), k(t) \in C^1([0, T])$, $w(t, x) \in C(\overline{D}_T) \cap C^{1,2}(D_T)$ *with* $D_T = \{(t, x) \in \mathbb{R}^2 \mid 0 < t \leqslant T,\ l(t) < x < k(t)\}$*, and*

$$\begin{cases} w_t \geqslant w_{xx} - \beta w_x + f(w), & 0 < t \leqslant T,\ l(t) < x < k(t), \\ w \geqslant u, & 0 < t \leqslant T,\ x = l(t), \\ w = 0, \quad k'(t) \geqslant -\mu w_x, & 0 < t \leqslant T,\ x = k(t), \end{cases}$$

with

$$g(t) \leqslant l(t) \leqslant h(t) \text{ for } t \in [0, T], \quad h_0 \leqslant k(0), \quad u_0(x) \leqslant w(0, x) \text{ for } x \in [l(0), h_0], \tag{19}$$

where (u, g, h) *is a solution of* (P)*. Then*

$$h(t) \leqslant k(t) \text{ for } t \in (0, T], \quad u(t, x) \leqslant w(t, x) \text{ for } t \in (0, T] \text{ and } l(t) < x < h(t).$$

The proof of Lemma 3.1 is identical to that of Lemma 5.7 in [12], and a minor modification of this proof yields Lemma 3.2.

Remark 3.3. The function $\bar{u}$, or the triple $(\bar{u}, \bar{g}, \bar{h})$ in Lemma 3.1 and the function w, or the triple (w, l, k) in Lemma 3.2 are often called the upper solutions of (P). There is a symmetric version of Lemma 3.2, where the conditions on the left and right boundaries are interchanged. The lower solutions can be defined analogously by reversing all the inequalities except for $g(t) \leqslant l(t) \leqslant h(t)$ in (19). We also have corresponding comparison results for lower solutions in each case.

3.2. Phase plane analysis and stationary solutions

We first use the phase plane analysis to study the following equation

$$q''(z) + \gamma q'(z) + f(q) = 0, \quad q(z) \geqslant 0 \quad \text{for } z \in J, \tag{20}$$

where J is some interval in $\mathbb{R}$. Note that a nonnegative stationary solution u of $(P)_1$ solves (20) with $\gamma = -\beta$, a nonnegative traveling wave $u(t, x) = q(x - ct)$ of $(P)_1$ solves (20) with $\gamma = c - \beta$ and $J = \mathbb{R}$.

The equation (20) is equivalent to the system

$$\begin{cases} q'(z) = p, \\ p'(z) = -\gamma p - f(q). \end{cases} \tag{21}$$



Fig. 1. Trajectories corresponding to the equation $q'' + \gamma q' + f(q) = 0$. (a) The case $\gamma \in (-c_0, c_0)$; (b) the case $\gamma \leqslant -c_0$.

A solution $(q(z), p(z))$ of this system traces out a trajectory in the q–p phase plane (cf. [2,13,31]). Such a trajectory has a slope

$$\frac{\mathrm{d}p}{\mathrm{d}q} = -\gamma - \frac{f(q)}{p} \tag{22}$$

at any point where $p \neq 0$. It is easily seen that $(0,0)$ and $(1,0)$ are two singular points on the phase plane. We are only interested in the case $\gamma < c_0 := 2\sqrt{f'(0)}$. For such a γ, the eigenvalues of the corresponding linearizations at the singular points are

$$\lambda_0^{\pm} = \frac{-\gamma \pm \sqrt{\gamma^2 - 4f'(0)}}{2} \quad (\text{at } (0,0)) \quad \text{and} \quad \lambda_1^{\pm} = \frac{-\gamma \pm \sqrt{\gamma^2 - 4f'(1)}}{2} \quad (\text{at } (1,0)),$$

respectively. Since $f'(0) > 0$ and $f'(1) < 0$, $(1,0)$ is a saddle point, $(0,0)$ is a center when $\gamma = 0$, or a focus when $0 < |\gamma| < c_0$, or a node when $\gamma \leqslant -c_0$. By the phase plane analysis (cf. [2,13,31]), it is not difficult to give all kinds of bounded, nonnegative solutions of (20) for $\gamma < c_0$ (see Fig. 1).

(i) **Constant solutions:** $q \equiv 0$ and $q \equiv 1$.

(ii) **Strictly decreasing solutions on the half-line in case** $\gamma < c_0$**:** $q(\cdot) = U(\cdot - z_0; \gamma)$ for any $z_0 \in \mathbb{R}$, where $U \in C^2((-\infty, 0])$ is the unique solution of (20) in $(-\infty, 0)$, with $U(0;\gamma) = 0$, $U(-\infty;\gamma) = 1$ and $U'(\cdot\,;\gamma) < 0$ in $(-\infty, 0]$ (see Γ_1 and Γ_5 in Fig. 1). Denote

$$P(\gamma) := -\mu U'(0;\gamma).$$

Using the comparison principle for the ordinary differential equation (22) we have $P'(\gamma) < 0$ for $\gamma \in (-\infty, c_0)$, $P(c_0 - 0) = 0$ and $P(-\infty) = +\infty$ (see Fig. 2(a)).

(iii) **Strictly increasing solutions on the half-line in case** $\gamma \in (-c_0, c_0)$**:** $q(\cdot) = U_l(\cdot - z_0; \gamma)$ for any $z_0 \in \mathbb{R}$, where $U_l \in C^2([0, \infty))$ is the unique solution of (20) in $(0, \infty)$, with $U_l(0;\gamma) = 0$, $U_l(\infty;\gamma) = 1$ and $U_l'(\cdot\,;\gamma) > 0$ in $[0, +\infty)$ (see Γ_4 in Fig. 1(a)).

(iv) **Solutions with compact supports in case** $\gamma \in (-c_0, c_0)$**:** $q(\cdot) = W(\cdot - z_0; b, \gamma)$ for any $z_0 \in \mathbb{R}$, where for each $b \in (0, P(\gamma))$, there exists a unique $L(b,\gamma) > 0$ such that $W \in C^2([-L(b,\gamma), 0])$ is the unique solution of (20) in $(-L(b,\gamma), 0)$



Fig. 2. (a) The γ–b plane about stationary solutions: each point in S_1 (resp. S_2) corresponds to a compactly supported solution (resp. a tadpole-like solution); (b) the c–b plane about traveling waves, point A corresponds to a rightward traveling semi-wave satisfying both the equation and Stefan boundary condition.

with $W(-L(b,\gamma);b,\gamma) = W(0;b,\gamma) = 0$ and $b = -\mu W'(0;b,\gamma)$ (see Γ_2 and Γ_3 in Fig. 1(a)). Each point (γ, b) in the set $S_1 := \{(\gamma,b) \mid 0 < b < P(\gamma),\ -c_0 < \gamma < c_0\}$ in Fig. 2(a) corresponds to such a compactly supported solution $W(z;b,\gamma)$.

(v) **Tadpole-like solutions in case** $\gamma \leqslant -c_0$**:** $q(\cdot) = V(\cdot - z_0;b,\gamma)$ for any $z_0 \in \mathbb{R}$, where for each $b \in (0, P(\gamma))$, $V \in C^2((-\infty,0])$ is the unique solution of (20) in $(-\infty,0)$ with $V(0;b,\gamma) = 0$, $V(-\infty;b,\gamma) = 0$ and $b = -\mu V'(0;b,\gamma)$ (see Γ_6 and Γ_7 in Fig. 1(b)). Each point (γ,b) in the set $S_2 := \{(\gamma,b) \mid 0 < b < P(\gamma),\ \gamma \leqslant -c_0\}$ in Fig. 2(a) corresponds to such a tadpole-like solution $V(z;b,\gamma)$. In addition, it is easily seen from the phase plane that there exists a unique $\bar{z} < 0$ such that $V(z;b,\gamma)$ is strictly increasing in $(-\infty,\bar{z})$ and strictly decreasing in $(\bar{z},0]$.
We call V a *tadpole-like solution* since its graph has a big "head" and a boundary on the right side, and an infinite long "tail" on the left side. Similarly, when we construct a traveling wave with the form $V(x-ct;b,\gamma)$, we call it a *tadpole-like traveling wave*.

(vi) **Strictly increasing solutions in** $\mathbb{R}$ **in case** $\gamma \leqslant -c_0$**:** $q(\cdot) = Q(\cdot - z_0;\gamma)$ for any $z_0 \in \mathbb{R}$, where $Q \in C^2(\mathbb{R})$ is the unique solution of (20) in $\mathbb{R}$ with $Q(-\infty;\gamma) = 0$, $Q(\infty;\gamma) = 1$, $Q(0;\gamma) = 1/2$ and $Q'(z;\gamma) > 0$ in $\mathbb{R}$ (see Γ_8 in Fig. 1(b)).

Each nonnegative stationary solution of $(P)_1$ is a solution of the problem (20) with $\gamma = -\beta$. Using the above results we see that, when $\beta \in (0,c_0)$, a bounded, nonnegative stationary solution of $(P)_1$ is either 0, or 1, or a strictly decreasing solutions $U(\cdot - z_0;-\beta)$ $(z_0 \in \mathbb{R})$ defined on $(-\infty,z_0]$, or a compactly supported solution $W(\cdot - z_0;b,-\beta)$ $(z_0 \in \mathbb{R})$ for some $b \in (0,P(-\beta)) = (0,-\mu U'(0;-\beta))$, or a strictly increasing solution $U_l(\cdot - z_0;-\beta)$ $(z_0 \in \mathbb{R})$ defined on $[z_0,\infty)$. When $\beta \geqslant c_0$, a bounded, nonnegative stationary solution of $(P)_1$ is either 0, or 1, or a strictly decreasing solution $U(\cdot - z_0;-\beta)$ $(z_0 \in \mathbb{R})$ defined on $(-\infty,z_0]$, or a tadpole-like function $V(\cdot - z_0;b,-\beta)$ $(z_0 \in \mathbb{R})$ for some $b \in (0,P(-\beta)) = (0,-\mu U'(0;-\beta))$, or a strictly increasing solution $Q(\cdot - z_0;-\beta)$ $(z_0 \in \mathbb{R})$ defined in $\mathbb{R}$.

3.3. Traveling waves

If $u(t,x) = q(x-ct)$ is a traveling wave of $u_t = u_{xx} - \beta u_x + f(u)$, then (c,q) solves (20) with $\gamma = c - \beta$, that is,



Fig. 3. Points A and F correspond to traveling semi-waves, points B, C, E correspond to tadpole-like traveling waves, point D corresponds to a compactly supported traveling wave, the waves denoted by A, B, C, D satisfy Stefan boundary condition. (a) The case $c_0 \leqslant \beta < \beta^*$; (b) the case $\beta = \beta^*$.

$$q''(z) + (c-\beta)q'(z) + f(q) = 0. \tag{23}$$

In this paper we will use several types of traveling waves which are specified now.

(I) Traveling wave $Q(x-ct; c-\beta)$ for any $c \leqslant \beta - c_0$, where $q(z) = Q(z; c-\beta)$ satisfies (23) and

$$q(-\infty) = 0,\ q(\infty) = 1,\ q(0) = \frac{1}{2},\ q'(z) > 0 \text{ for } z \in \mathbb{R}. \tag{24}$$

The existence of such solutions has been given in the previous subsection.

(II) Rightward traveling semi-wave $U^*(x - c^*t)$ with $c^* \in (0, c_0 + \beta)$, where $q(z) = U^*(z) := U(z; c^* - \beta)$ satisfies (23) with $c = c^*$ and

$$q(0) = 0,\ q(-\infty) = 1,\ -\mu q'(0) = c^*,\ q'(z) < 0 \text{ for } z \in (-\infty, 0], \tag{25}$$

that is, $(c^*, U^*(z))$ is a solution of (5) (cf. point A in Fig. 2(b) and in Fig. 3). $U^*(x-c^*t)$ is called a traveling semi-wave as in [13] since $U^*(z)$ is defined only on the half-line $(-\infty, 0]$.

Lemma 3.4. *Assume $\beta > 0$. Then*

(i) *there exists a unique $c^* = c^*(\beta) \in (0, c_0 + \beta)$ such that the problems (23) and (25) with $c = c^*$ have a solution, which is unique and denoted by $U^*(z)$;*
(ii) $0 < \frac{\mathrm{d}}{\mathrm{d}\beta} c^*(\beta) < 1$ *for $\beta > 0$;*
(iii) *there exists a unique $\beta^* > c_0$ such that*

$$c^*(\beta) - \beta + c_0 > 0 \ \ (\textit{resp.}\ = 0, < 0) \textit{ when } \beta < \beta^* \ \ (\textit{resp. } \beta = \beta^*,\ \beta > \beta^*). \tag{26}$$

Proof. (i) For any $c < c_0 + \beta$, the problem (20) with $\gamma = c - \beta < c_0$ has a unique strictly decreasing solution $q(\cdot) = U(\cdot\,; c-\beta)$ in $(-\infty, 0]$, satisfying $U(0; c-\beta) = 0$, $U(-\infty; c-\beta) = 1$ and $U'(\cdot\,; c-\beta) < 0$ in $(-\infty, 0]$. Denote $P(c-\beta) := -\mu U'(0; c-\beta)$ as above, then $P(c-\beta)$ is strictly decreasing in $c \in (-\infty, c_0 + \beta)$,

$$(P(c-\beta)-c)\big|_{c=0} = P(-\beta) > 0 \quad \text{and} \quad (P(c-\beta)-c)\big|_{c=c_0+\beta-0} = -c_0-\beta < 0$$

(see Fig. 2(b)). Hence the equation $P(c-\beta) = c$ has a unique root $c = c^*(\beta) \in (0, c_0+\beta)$, that is,

$$c^*(\beta) = P(c^*(\beta)-\beta) = -\mu U'(0; c^*(\beta)-\beta). \tag{27}$$

(ii) Differentiating $P(c^*(\beta)-\beta) = c^*(\beta)$ in β and using the fact $P'(\gamma) < 0$ for $\gamma < c_0$ we have

$$\frac{\mathrm{d}c^*(\beta)}{\mathrm{d}\beta} = \frac{-P'(c^*(\beta)-\beta)}{1-P'(c^*(\beta)-\beta)} \in (0,1).$$

(iii) Set $\beta^* := P(-c_0)+c_0 > c_0$. Then $c = \beta^*-c_0$ is a root of $P(c-\beta^*) = c$ in $(0, c_0+\beta^*)$. By the definition of $c^*(\beta)$ and by its uniqueness we have $\beta^*-c_0 = c^*(\beta^*)$. Moreover, the inequalities in (ii) show that the function $c^*(\beta)-\beta+c_0$ is strictly decreasing in $\beta > 0$ and so it has a unique zero β^*. This proves (26). □

(III) Leftward traveling semi-wave $U_l^*(x-c_l^*t)$ in case $\beta \in (0, c_0)$, where $c_l^* = c_l^*(\beta) \in (\beta-c_0, 0)$, $q(z) = U_l^*(z) := U_l(z; c_l^*-\beta)$ satisfies (23) with $c = c_l^*$ and

$$q(0) = 0,\ q(\infty) = 1,\ -\mu q'(0) = c_l^*,\ q'(z) > 0 \text{ for } z \in [0,\infty). \tag{28}$$

For any given $\beta \in (0, c_0)$, the existence and uniqueness of such a solution can be proved as in Lemma 3.4(i).

(IV) Tadpole-like traveling wave $V(x-ct; b, c-\beta)$ in case $\beta > c_0$. For any $c \in (0, \beta-c_0]$ and any $b \in (0, P(c-\beta))$, $V(x-ct; b, c-\beta)$ is a tadpole-like traveling wave if the function $q(z) := V(z; b, c-\beta)$ satisfies (23) and

$$q(0) = q(-\infty) = 0,\ q(z) > 0 \text{ for } z \in (-\infty, 0) \text{ and } -\mu q'(0) = b \tag{29}$$

(cf. points B, C, E in Fig. 3(a)). In particular, when $b = c$, the function $V(z; c, c-\beta)$ is a solution of (23) and

$$q(0) = q(-\infty) = 0,\ q(z) > 0 \text{ for } z \in (-\infty, 0) \text{ and } -\mu q'(0) = c \tag{30}$$

(cf. points B, C in Fig. 3(a)). On the existence of such solutions we have the following results.

Lemma 3.5. *Let β^* be the constant given in Lemma 3.4. Assume $c_0 < \beta < \beta^*$. Then*

(i) *for any $b \in (0, P(-c_0))$, (29) and (23) with $c = \beta - c_0$ have a unique tadpole-like solution $V(z; b, -c_0)$ (cf. points B, E in Fig. 3(a)). Moreover, there exists $z_b < 0$*

such that

$$V(\cdot + z_b; b, -c_0) \to Q(\cdot) \text{ locally uniformly in } \mathbb{R}, \quad \text{as } b \to P(-c_0); \tag{31}$$

(ii) $q(z) = V^*(z) := V(z; \beta - c_0, -c_0)$ *is the unique tadpole-like solution of* (23) *and* (30) *with* $c = \beta - c_0$, *that is, the unique solution of* (13);

(iii) *for any* $\delta \in (0, \beta - c_0)$, $q(z) = V_\delta(z) := V(z; \beta - c_0 - \delta, -c_0 - \delta)$ *is a tadpole-like solution of* (23) *and* (30) *with* $c = \beta - c_0 - \delta$. *Moreover,* $V_\delta(z) \to V^*(z)$ *locally uniformly in* $(-\infty, 0]$ *as* $\delta \to 0$ *(cf. point* C *in Fig. 3(a)).*

Proof. (i) Since $c_0 < \beta < \beta^*$, we have $0 < \beta - c_0 < c^*$ by Lemma 3.4. On the c–b plane (see Fig. 3(a)), any point $E(\beta - c_0, b)$ with $b \in (0, P(-c_0))$ corresponds to a tadpole-like solution $V(z; b, -c_0)$ of (29) and (23) with $c = \beta - c_0$ (cf. trajectories Γ_6 or Γ_7 in Fig. 1(b). Note that such a solution does not necessarily satisfy Stefan condition since b may be not equal to c). As $b \to P(\beta - c_0)$ (i.e., point E moves up to F in Fig. 3(a)), the trajectory of $V(z; b, -c_0)$ approaches the union of the trajectories of Q and U^* (i.e., $\Gamma_6 \to \Gamma_5 \cup \Gamma_8$). Denote $z_b := \min\{z < 0 \mid V(z; b, -c_0) = \frac{1}{2}\}$. Then the trajectory of $V(\cdot + z_b; b, -c_0)$ approaches that of $Q(\cdot)$ (since $Q(0) = \frac{1}{2}$), and so we obtain (31) by continuity.

(ii) On the c–b plane, the line $\{b = c\}$ passes through the domain $S_2(\beta) := \{(c, b) \mid 0 < b < P(c - \beta),\ 0 < c < \beta - c_0\}$ and leaves it at a point $B(\beta - c_0, \beta - c_0)$. This point corresponds to the desired tadpole-like solution $V^*(z) := V(z; \beta - c_0, -c_0)$.

(iii) For any small $\delta > 0$, we consider the point $C(\beta - c_0 - \delta, \beta - c_0 - \delta)$ on the c–b plane. Since $C \in S_2(\beta) \cap \{b = c\}$, it corresponds to a tadpole-like solution $V_\delta(z) := V(z; \beta - c_0 - \delta, -c_0 - \delta)$ of (23) and (30) with $c = \beta - c_0 - \delta$. In particular, $V_\delta(z)$ satisfies the following initial value problem

$$\begin{cases} q''(z) - (c_0 + \delta)q'(z) + f(q) = 0, & z < 0, \\ q(0) = 0, \quad -\mu q'(0) = \beta - c_0 - \delta. \end{cases}$$

Since V^* satisfies this problem with $\delta = 0$ and since V_δ depends on δ continuously, we have $V_\delta(\cdot) \to V^*(\cdot)$ as $\delta \to 0$, uniformly in $[-M, 0]$ for any $M > 0$. This proves the lemma. □

In a similar way, one can prove the following lemma (cf. Fig. 3(b)).

Lemma 3.6. *Assume* $\beta = \beta^*$. *Then for any small* $\delta > 0$, $q(z) = V_\delta^*(z) := V(z; \beta^* - c_0 - \delta, -c_0 - \delta)$ *is the unique tadpole-like solution of* (23) *and* (30) *with* $\beta = \beta^*$, $c = \beta^* - c_0 - \delta$. *Moreover,* $V_\delta^*(\cdot) \to U^*(\cdot)$ *locally uniformly in* $(-\infty, 0]$ *as* $\delta \to 0$, *where* $U^*(z)$ *is the unique solution of* (23) *and* (25) *with* $\beta = \beta^*$, $c^* = c^*(\beta^*) = \beta^* - c_0$.

(V) Compactly supported traveling wave $W(x - ct; c, c - \beta)$ in case $\beta \in [c_0, \beta^*)$, where for any $c \in (\beta - c_0, c^*(\beta))$, $q(z) = W(z; c, c - \beta)$ satisfies (23) and

$$q(0) = q(-L(c, c - \beta)) = 0, \quad q(z) > 0 \text{ in } (-L(c, c - \beta), 0) \quad \text{and} \quad -\mu q'(0) = c. \tag{32}$$

Lemma 3.7. *Assume $c_0 \leqslant \beta < \beta^*$. For any $\delta \in (0, c^*(\beta) - \beta + c_0)$, $q(z) = W_\delta(z) := W(z; \beta - c_0 + \delta, -c_0 + \delta)$ is the unique solution of the problem (23) and (32) with $c = \beta - c_0 + \delta$. Moreover, $L_\delta \to \infty$ and $D_\delta \to 1$ as $\delta \to c^*(\beta) - \beta + c_0$, where $L_\delta := L(\beta - c_0 + \delta, -c_0 + \delta)$ denotes the width of the support of $W_\delta(\cdot)$, and D_δ denotes its height.*

Proof. We only prove the case $\beta \in (c_0, \beta^*)$, the proof for the case $\beta = c_0$ is similar.

When $c_0 < \beta < \beta^*$, we have $c^*(\beta) - \beta + c_0 > 0$ by Lemma 3.4. On the c–b plane (see Fig. 3(a)), the line $b = c$ leaves the domain $S_2(\beta)$ and enters $S_1(\beta) := \{(c, b) \mid 0 < b < P(c - \beta),\ \beta - c_0 < c < \beta + c_0\}$ at a point $B(\beta - c_0, \beta - c_0)$, then it passes through $S_1(\beta)$ and leaves it finally at $A(c^*(\beta), c^*(\beta))$. For any $\delta \in (0, c^*(\beta) - \beta + c_0)$, the point $(c, b) = (\beta - c_0 + \delta, \beta - c_0 + \delta)$ is on the line segment AB (cf. point D in Fig. 3(a)), it corresponds to a trajectory like Γ_2 on the q–p phase plane, and so it defines a compactly supported function $W_\delta(z) := W(z; \beta - c_0 + \delta, -c_0 + \delta)$. As $\delta \to c^*(\beta) - \beta + c_0$, the point $(\beta - c_0 + \delta, \beta - c_0 + \delta)$ approaches point A in Fig. 3(a), this implies that its corresponding trajectory approaches the union of the trajectories of $U^*(\cdot)$ and $U_l(\cdot; c^*(\beta) - \beta)$ (i.e., $\Gamma_2 \to \Gamma_1 \cup \Gamma_4$ in Fig. 1(a)). Therefore, the corresponding function W_δ satisfies $\max\limits_{-L_\delta \leqslant z \leqslant 0} W_\delta(z) = D_\delta \to 1$, and the width L_δ of its support tends to ∞ as $\delta \to c^*(\beta) - \beta + c_0$. $\square$

3.4. Zero number arguments

In what follows, we use $\mathcal{Z}_I[w(\cdot)]$ to denote the number of zeros of a continuous function $w(\cdot)$ defined in $I \subset \mathbb{R}$. The following lemma is an easy consequence of the proofs of Theorems C and D in Angenent [1].

Lemma 3.8. *(See [14, Lemma 2.1].) Let $u : [0, T] \times [0, 1] \to \mathbb{R}$ be a bounded classical solution of*

$$u_t = a(t, x)u_{xx} + b(t, x)u_x + c(t, x)u \tag{33}$$

with boundary conditions

$$u(t, 0) = l_0(t),\ \ u(t, 1) = l_1(t),$$

where $l_0, l_1 \in C^1([0, T])$, and each function is either identically zero or never zero for $t \in [0, T]$. In the special case where $l_0(t), l_1(t) \equiv 0$ we assume further that $u(t, \cdot) \not\equiv 0$ for each $t \in [0, T]$. Suppose also that

$$a, 1/a, a_t, a_x, a_{xx}, b, b_t, b_x, c \in L^\infty,\ \ \text{and } u(0, \cdot) \not\equiv 0 \text{ when } l_0 = l_1 \equiv 0.$$

Then for each $t \in (0,T]$, $\mathcal{Z}_{[0,1]}[u(t,\cdot)] < \infty$. Moreover, $\mathcal{Z}_{[0,1]}[u(t,\cdot)]$ is nonincreasing in t for $t \in (0,T]$, and if for some $t_0 \in (0,T]$ the function $u(t_0,\cdot)$ has a degenerate zero $x_0 \in [0,1]$, then $\mathcal{Z}_{[0,1]}[u(t_1,\cdot)] > \mathcal{Z}_{[0,1]}[u(t_2,\cdot)]$ for all $t_1, t_2 \in (0,T]$ satisfying $t_1 < t_0 < t_2$.

The following variant of Lemma 3.8 is convenient for our applications.

Lemma 3.9. *(See [14, Lemma 2.2].) Let $\xi_1(t) < \xi_2(t)$ be two continuous functions for $t \in (t_0,t_1)$. If $u(t,x)$ is a continuous function for $t \in (t_0,t_1)$ and $x \in J(t) := [\xi_1(t), \xi_2(t)]$, and satisfies (33) in the classical sense for such (t,x), with*

$$u(t,\xi_1(t)) \neq 0,\ \ u(t,\xi_2(t)) \neq 0 \text{ for } t \in (t_0,t_1),$$

then for each $t \in (t_0,t_1)$, $\mathcal{Z}_{J(t)}[u(t,\cdot)] < \infty$. Moreover $\mathcal{Z}_{J(t)}[u(t,\cdot)]$ is nonincreasing in t for $t \in (t_0,t_1)$, and if for some $s \in (t_0,t_1)$ the function $u(s,\cdot)$ has a degenerate zero $x_0 \in J(s)$, then $\mathcal{Z}_{J(s_1)}[u(s_1,\cdot)] > \mathcal{Z}_{J(s_2)}[u(s_2,\cdot)]$ for all s_1, s_2 satisfying $t_0 < s_1 < s < s_2 < t_1$.

In our approach we will compare the solution u of (P) with traveling semi-wave U^* or tadpole-like traveling wave V^* or compactly supported traveling wave W_δ by studying the number of their intersection points. Now we give some preliminary results.

We use $\Psi(x-ct-C)$ (for some $c>0$ and some $C \in \mathbb{R}$) to represent one of the traveling waves $U^*(x-c^*t-C)$, $V^*(x-(\beta-c_0)t-C)$ and $W_\delta(x-(\beta-c_0+\delta)t-C)$. Denote the support of $\Psi(x-ct-C)$ by $[k_1(t),k_2(t)]$, where $k_2(t)=ct+C$, $k_1(t)=-\infty$ in case $\Psi=U^*$ or $\Psi=V^*$, and $k_1(t)=k_2(t)-L_\delta$ in case $\Psi=W_\delta$. Denote

$$r(t) := \min\{h(t),k_2(t)\}, \quad R(t) := \max\{h(t),k_2(t)\}, \quad l(t) := \max\{g(t),k_1(t)\}$$

and

$$\eta(t,x) := u(t,x) - \Psi(x-ct-C), \quad x \in J(t) := [l(t),r(t)],\ t \in (t_1,t_2)$$

(here we only consider the case where $J(t) \neq \emptyset$ for each $t \in (t_1,t_2)$, otherwise, u and Ψ have no common domain and so there is no need to compare them). We notice that η satisfies

$$\eta_t = \eta_{xx} - \beta\eta_x + c(t,x)\eta \quad \text{for } x \in (l(t),r(t)),\ t \in (t_1,t_2)$$

with $c(t,x) := [f(u(t,x)) - f(\Psi(x-ct-C))]/\eta(t,x)$ when $\eta(t,x) \neq 0$, and $c(t,x)=0$ otherwise. Using Lemmas 3.8 and 3.9 one can obtain the following result on the number of zeros of $\eta(t,\cdot)$.

Lemma 3.10. *For any given $C \in \mathbb{R}$, let $r(t)$, $R(t)$, $l(t)$ and η be defined as above.*

(a) *Assume that* $r(t_1) < R(t_1)$ *and* $k_1(t) \neq g(t)$ *for all* $t \in (t_1, t_2)$. *Then*
  (i) $\mathcal{Z}_{J(t)}[\eta(t,\cdot)]$ *is finite and nonincreasing in* $t \in (t_1, t_2)$;
  (ii) if there exists $t_0 \in (t_1, t_2)$ *such that* $r(t_0) = R(t_0)$, *or* $\eta(t_0, \cdot)$ *has a degenerate zero in the interior of* $J(t_0)$, *then* $\mathcal{Z}_{J(\tau_1)}[\eta(\tau_1,\cdot)] > \mathcal{Z}_{J(\tau_2)}[\eta(\tau_2,\cdot)]$ *for any* $t_1 < \tau_1 < t_0 < \tau_2 < t_2$.

(b) *Assume that* $\Psi = W_\delta$, $h(t_1) - g(t_1) > L_\delta$, $\mathcal{Z}_{J(t_1)}[\eta(t_1,\cdot)] = 1$. *Assume also that* $k_1(t_*) = g(t_*)$ *for some* $t_* \in (t_1, t_2)$. *Then* $\mathcal{Z}_{J(t)}[\eta(t,\cdot)] = 1$ *for* $t \in (t_1, t_*)$ *and there exists* $\tilde{s} \in (t_*, t_2)$ *such that, when* $t \in (t_*, \tilde{s})$, *either* $\mathcal{Z}_{J(t)}[\eta(t,\cdot)] = 0$ *(and so* $W_\delta(x - ct - C) < u(t,x)$ *in* $[k_1(t), k_2(t)]$*), or* $\mathcal{Z}_{J(t)}[\eta(t,\cdot)] = 2$.

Proof. The proof is based on [18, Theorem 2] and is essentially identical to that of Lemma 2.3 in [14]. Hence we only give a sketch here and omit the details.

(a) Since $r(t_1) < R(t_1)$, we have $h(t) \neq k_2(t)$ for $t_1 < t \ll t_1 + 1$. Let s_1 be the first time in (t_1, t_2) such that $r(t) = R(t)$, that is, $s_1 := \sup\{s \in (t_1, t_2) \mid r(t) < R(t) \text{ for } t \in (t_1, s)\}$. If $s_1 = t_2$, then the conclusions in (a) follow from Lemma 3.9 directly. In what follows we assume that $s_1 \in (t_1, t_2)$. By the definition of s_1 we see that $\eta(t, r(t)) \neq 0$ for all $t \in (t_1, s_1)$. On the other hand $\eta(t, l(t)) \neq 0$ by the assumption $k_1(t) \neq g(t)$ for $t \in (t_1, t_2)$. By Lemma 3.9, $\mathcal{Z}_{J(t)}[\eta(t,\cdot)]$ is finite and nondecreasing in $t \in (t_1, s_1)$, and each time a degenerate zero appears in the interior of $J(t)$ for $\eta(t,\cdot)$ the value of $\mathcal{Z}_{J(t)}[\eta(t,\cdot)]$ is decreased by at least 1. These facts imply that in the interval (t_1, s_1) there can exist at most finitely many values of t such that $\eta(t,\cdot)$ has a degenerate zero. Thus we can find $\epsilon \in (0, s_1 - t_1)$ such that for $t \in [s_1 - \epsilon, s_1)$, $\eta(t,\cdot)$ has only non-degenerate zeros in $J(t)$. Due to the non-degeneracy, the zeros of $\eta(t,\cdot)$, with $t \in [s_1 - \epsilon, s_1)$, can be expressed as smooth curves:

$$l(t) < z_1(t) < z_2(t) < \cdots < z_m(t) < r(t), \quad t \in [s_1 - \epsilon, s_1).$$

By [18, Theorem 2] one can prove that $\lim_{t \to s_1 - 0} z_i(t)$ exists (denoted by $\bar{z}_i$) and $\{\bar{z}_i\}_{i=1}^m$ are the only zeros of $\eta(s_1,\cdot)$ (maybe $\bar{z}_i = \bar{z}_{i+1}$ for some i). Moreover, by the maximum principle we have $\bar{z}_m = r(s_1)$, that is, the largest zero $z_m(t)$ tends to the right boundary. Then using the maximum principle again in the domain $\{(t,x) \mid s_1 < t < s_1 + \epsilon_1,\ r(t) - \epsilon_1 < x < r(t)\}$ for some small $\epsilon_1 > 0$, we can show that the boundary zero $r(s_1)$ disappears immediately after time s_1. In summary,

$$\mathcal{Z}_{J(\tau_1)}[\eta(\tau_1,\cdot)] = m \geqslant \mathcal{Z}_{J(s_1)}[\eta(s_1,\cdot)] > \mathcal{Z}_{J(\tau_2)}[\eta(\tau_2,\cdot)] \tag{34}$$

for $s_1 - \epsilon < \tau_1 < s_1 < \tau_2 < s_1 + \epsilon_1$. This conclusion implies that each time a zero for $\eta(t,\cdot)$ appears on the right boundary (which must be a degenerate one by the Stefan boundary condition), the value of $\mathcal{Z}_{J(t)}[\eta(t,\cdot)]$ is decreased by at least 1. Therefore, there can exist at most finitely many values $s_1, s_2, \cdots, s_n \in (t_1, t_2)$ such that, for each $i \in \{1, 2, \cdots, n\}$, $\eta(s_i, r(s_i)) = 0$, that is, $r(s_i) = R(s_i)$. Clearly, (34) holds with s_1 replaced by each s_i. Finally, using Lemma 3.9 in the intervals $(t_1, s_1), (s_1, s_2), \cdots, (s_{n-1}, s_n), (s_n, t_2)$ and using (34) at each time s_i we obtain the desired conclusions.

(b) Since $k_1'(t) = c > 0$ and $g'(t) < 0$ we have $k_1(t) < g(t)$ for $t \in (t_1, t_*)$ and $k_1(t) > g(t)$ for $t \in (t_*, t_2)$. Hence

$$\begin{aligned} &\eta(t, l(t)) = \eta(t, g(t)) < 0 \text{ for } t \in (t_1, t_*), \\ &\eta(t, l(t)) = \eta(t, k_1(t)) > 0 \text{ for } t \in (t_*, t_2). \end{aligned} \tag{35}$$

The assumption $h(t_1) - g(t_1) > L_\delta$ implies that $h(t) > g(t) + L_\delta \geqslant k_1(t) + L_\delta = k_2(t)$ for $t \in (t_1, t_*]$. Hence there exists $\hat{s} \in (t_*, t_2)$ such that $h(t) > k_2(t)$, and so

$$\eta(t, r(t)) = \eta(t, k_2(t)) > 0 \text{ for } t \in (t_1, \hat{s}). \tag{36}$$

Combining with (35) we see that $\eta(t, \cdot)$ has at least one zero in $(l(t), r(t))$ in the time interval (t_1, t_*). On the other hand, the assumption $\mathcal{Z}_{J(t_1)}[\eta(t_1, \cdot)] = 1$ and Lemma 3.9 imply that the zero is unique (denoted by $\gamma(t)$). This proves $\mathcal{Z}_{J(t)}[\eta(t, \cdot)] = 1$ for $t \in (t_1, t_*)$.

Using [18, Theorem 2] as above one can see that $\bar{z} = \lim_{t \to t_* - 0} \gamma(t) \in [l(t_*), r(t_*)]$ exists. $\bar{z} \neq r(t_*)$ due to (36). In case $\bar{z} = l(t_*)$, it is the unique zero of $\eta(t_*, \cdot)$ in $J(t_*)$ and $\eta(t_*, x) > 0$ for $x \in (\bar{z}, r(t_*)]$. Combining with (35) and (36), and using the maximum principle in the domain $\{(t, x) \mid t_* < t < \hat{s},\ l(t) < x < r(t)\}$, we deduce that $\eta(t, x) > 0$ in this domain, and so $\mathcal{Z}_{J(t)}[\eta(t, \cdot)] = 0$ for $t \in (t_*, \hat{s})$. This implies that $W_\delta(x - ct - C) < u(t, x)$ for $t \in (t_*, \hat{s})$ and $x \in [k_1(t), k_2(t)] \subset (g(t), h(t))$. In case $\bar{z} \in (l(t_*), r(t_*))$, $\eta(t_*, \cdot)$ has exactly two zeros $l(t_*)$ and $\bar{z}$ in $J(t_*)$. Moreover, $\eta(t_*, x) < 0$ for $x \in (l(t_*), \bar{z})$ and $\eta(t_*, x) > 0$ for $x \in (\bar{z}, r(t_*)]$. Set $z_* := (l(t_*) + \bar{z})/2$, then there exists $\tilde{s} \in (t_*, \hat{s})$ such that $\eta(t, z_*) < 0$ for $t \in [t_*, \tilde{s}]$. Combining with (36) and the second inequality in (35) we see that, when $t \in (t_*, \tilde{s})$, $\eta(t, \cdot)$ has zeros in $(l(t), z_*)$ and in $(z_*, r(t))$. It follows from the maximum principle that the zero is unique in $(l(t), z_*)$. It follows from Lemma 3.9 that the zero is also unique in $(z_*, r(t))$. Consequently we have $\mathcal{Z}_{J(t)}[\eta(t, \cdot)] = 2$ for $t \in [t_*, \tilde{s})$. This proves the second case in (b). $\square$

As can be expected, the presence of the advection makes the maximum points prefer to move rightward. Indeed we can show that the local maximum points concentrate near the right boundary under certain conditions.

Using zero number properties Lemma 3.9 to u_x, we see that $u_x(t, \cdot)$ has only non-degenerate zeros for all large t. Hence, $u(t, \cdot)$ has fixed number of (non-degenerate) local maximum points for large t.

Lemma 3.11. *Assume, for some $T \geqslant 0$, $u(t, \cdot)$ has exactly N (N is a positive integer) local maximum points $\{\xi_i(t)\}_{i=1}^N$ for all $t \geqslant T$, with*

$$g(t) < \xi_1(t) < \xi_2(t) < \cdots < \xi_N(t) < h(t).$$

If $N \geqslant 2$, then

$$\xi_1(t) \geqslant \rho(t) := \beta \cdot (t - T) + \frac{1}{2}\left[g(T) + \xi_1(T)\right] \quad \textit{for } T \leqslant t < T_\infty, \tag{37}$$

where $T_\infty = \sup\{\tau > T \mid \rho(t) < h(t) \textit{ in } [T, \tau]\}$.

Proof. Set $T_1 := \sup\{\tau \in [T, T_\infty) \mid \rho(t) < \xi_1(s) \text{ in } [T, \tau]\}$. Since $\rho(T) = \frac{1}{2}\left[g(T) + \xi_1(T)\right] < \xi_1(T)$ we have $T_1 > T$. If $T_1 = T_\infty$, then (37) holds. If $T < T_1 < T_\infty$, then we can derive a contradiction as follows. By the definitions of $\xi_1(t)$ and T_1 we have

$$u_x(t, x) > 0 \quad \text{for } x \in [g(t), \rho(t)],\ T \leqslant t < T_1, \tag{38}$$

and

$$u_x(T_1, \rho(T_1)) = u_x(T_1, \xi_1(T_1)) = 0. \tag{39}$$

Define

$$\zeta(t, x) := u(t, x) - u(t, 2\rho(t) - x) \quad \text{for } x \in [l(t), \rho(t)],\ t \geqslant T,$$

with $l(t) := \max\{g(t), 2\rho(t) - \xi_N(t)\}$. A direct calculation shows that

$$\zeta_t = \zeta_{xx} - \beta\zeta_x + c\zeta \quad \text{for } x \in [l(t), \rho(t)],\ t \in [T, T_1),$$

where c is a bounded function. It is easily seen that

$$\begin{cases} \zeta(T, x) < 0 & \text{for } x \in [l(T), \rho(T)], \\ \zeta(t, \rho(t)) = 0 & \text{for } t \in [T, T_1], \\ \zeta(t, l(t)) < 0 & \text{for } t \in [T, T_1]. \end{cases}$$

The last inequality follows from the following analysis. By the monotonicity of u_x and the fact that $\xi_N(t)$ is the rightmost local maximum point, we see that, if $\zeta(t_1, l(t_1)) = 0$ for some $t_1 \in (T, T_1]$ (denote by t_1 the first of such times), then $\zeta(t_1, l(t_1) + \varepsilon) > 0$ for some small $\varepsilon > 0$. This, however, is impossible since $\zeta(t_1, x) < 0$ for all $x \in (l(t_1), \rho(t_1))$ by the maximum principle and by the fact that $\zeta(t, l(t)) < 0$ for $t \in [T, t_1)$.

Now we use the Hopf lemma for ζ in the domain $\{(t, x) \mid l(t) \leqslant x \leqslant \rho(t), T \leqslant t \leqslant T_1\}$ to derive

$$\zeta(T_1, \rho(T_1)) = 0 \quad \text{and} \quad \zeta_x(T_1, \rho(T_1)) > 0.$$

The latter, however, contradicts (39). This proves $T_1 = T_\infty$. $\square$

3.5. Upper bound estimate

In order to study the convergence of u we give some precise upper bound estimates for the solutions. In this paper we always write

$$A := 2\max\left\{1, \|u_0\|_{L^\infty([-h_0,h_0])}\right\}. \tag{40}$$

1. *Bound of u near the free boundary $x = h(t)$.* For any $\delta \in (0, -f'(1))$, set

$$\bar{g}(t) := g(t), \quad \bar{h}(t) := c^* t - Me^{-\delta t} + H \quad \text{for some } M,\ H > 0,$$

and

$$\bar{u}(t,x) := (1 + Ae^{-\delta t})U^*(x - \bar{h}(t)) \quad \text{for } x \leqslant \bar{h}(t),\ t > 0,$$

where U^* is the rightward traveling semi-wave (the solution of (5)). A direct calculation as in the proof of [16, Lemma 3.2] shows that $(\bar{u}, \bar{g}, \bar{h})$ is an upper solution of (P) provided $M, H > 0$ are large. Hence we have

$$u(t,x) \leqslant \bar{u}(t,x) \text{ for } x \in [g(t), h(t)] \text{ and } t > 0, \quad h(t) \leqslant \bar{h}(t) \text{ for } t > 0. \tag{41}$$

2. *Bound of u in case $\beta \geqslant c_0$.* We define a function $f_A(s) \in C^2([0,\infty), \mathbb{R})$ such that

$$f_A(s) \begin{cases} = f'(0)s, & 0 \leqslant s \leqslant 1, \\ > 0, & 1 < s < A, \\ < 0, & s > A, \end{cases}$$
$$f_A'(A) < 0, \quad f(s) \leqslant f_A(s) \leqslant f'(0)s \text{ for } s \geqslant 0. \tag{42}$$

Denote by $Q_A(z)$ the unique solution of (4) with $c = c_0$, with f replaced by f_A and $q(+\infty) = 1$ replaced by $q(+\infty) = A$. Then by the comparison principle we have

$$u(t,x) \leqslant Q_A(x - (\beta - c_0)t + x_0) \quad \text{for } x \in [g(t), h(t)],\ t > 0, \tag{43}$$

provided $x_0 > 0$ is large enough.

Using f_A we consider the Cauchy problem:

$$\begin{cases} (u_1)_t = (u_1)_{xx} - \beta(u_1)_x + f_A(u_1), & x \in \mathbb{R},\ t > 0, \\ u_1(0,x) = \tilde{u}_0(x) := \begin{cases} u_0(x), & x \in [-h_0, h_0], \\ 0, & |x| > h_0. \end{cases} \end{cases}$$

Since $f_A \geqslant f$, by the comparison principle we have

$$u(t,x) \leqslant u_1(t,x), \quad x \in [g(t), h(t)],\ t > 0. \tag{44}$$

Set $y := x - \beta t$ and $u_2(t,y) := u_1(t,x) = u_1(t, y+\beta t)$. Then u_2 is a solution of

$$\begin{cases} (u_2)_t = (u_2)_{yy} + f_A(u_2), & y \in \mathbb{R},\ t > 0, \\ u_2(0,y) = \tilde{u}_0(y) \in [0,A], & y \in \mathbb{R}. \end{cases}$$

Set $\tilde{u}(t,y) := \frac{1}{A} u_2(t,-y)$, then $\tilde{u}(t,y)$ is the solution of

$$\begin{cases} \tilde{u}_t = \tilde{u}_{yy} + \frac{1}{A} f_A\big(A\tilde{u}\big), & y \in \mathbb{R},\ t > 0, \\ \tilde{u}(0,y) = \frac{1}{A}\tilde{u}_0(-y) \in [0,1], & y \in \mathbb{R}. \end{cases}$$

By (44) and by the definitions of u_2 and $\tilde{u}$ we have

$$u(t,x) \leqslant A\tilde{u}(t, \beta t - x), \quad x \in [g(t), h(t)],\ t > 0. \tag{45}$$

On the other hand, by Proposition 2.3 in [21] and its proof, there exist C_1, C_2, t_0 depending on u_0 such that

$$\tilde{u}\Big(t, c_0 t - \frac{3}{c_0}\ln\big(1 + \frac{t}{t_0}\big) + y\Big) \leqslant C_1 Z(t,y), \quad y \geqslant h_0,\ t > 0,$$

where

$$Z(t,y) := \frac{1}{\sqrt{t_0}} y e^{-\frac{c_0}{2}y}\big[C_2 e^{\frac{-y^2}{4(t+t_0)}} + h(t,y)\big], \quad y \in \mathbb{R},\ t > 0, \tag{46}$$

with $h(t,y)$ satisfying

$$\limsup_{t\to\infty} \sup_{0 \leqslant y \leqslant \sqrt{t+1}} |h(t,y)| \leqslant \frac{C_2}{2}.$$

In particular, there exists $C_3 > 0$ such that

$$\tilde{u}\Big(t, c_0 t - \frac{3}{c_0}\ln\big(1 + \frac{t}{t_0}\big) + y\Big) \leqslant C_3 e^{-\frac{5c_0}{12}y}, \quad y \in [0, \sqrt{t+1}\,].$$

Combining with (45) we have

$$u(t,x) \leqslant C_4 e^{-\frac{5c_0}{12}(Y(t)-x)} \quad \text{for } Y(t) - \sqrt{t+1} \leqslant x \leqslant \min\big\{Y(t), h(t)\big\},\ t \gg 1, \tag{47}$$

where $C_4 > 0$ is a constant and

$$Y(t) := (\beta - c_0)t + \frac{3}{c_0}\ln\big(1 + \frac{t}{t_0}\big), \quad t > 0. \tag{48}$$

Corollary 3.12.

(i) *Assume $\beta = c_0$. Then there exists C depending on u_0, c_0, h_0 such that*

$$u(t, -h_0) \leqslant Ct^{-\frac{5}{4}} \quad \textit{for } t > 0 \textit{ large.}$$

(ii) *Assume $\beta \in (c_0, \beta^*]$ and $h(t) = (\beta - c_0)t + O(1)$. Then there exists C depending on u_0, c_0, h_0 such that*

$$u(t, x) \leqslant Ct^{-\frac{5}{4}} \quad \textit{for } x \in \left[h(t) - \frac{\pi}{2}, h(t)\right] \textit{ and } t > 0 \textit{ large.}$$

Proof. We only prove (ii) since (i) can be proved similarly. In case (ii),

$$Y(t) = (\beta - c_0)t + \frac{3}{c_0} \ln\left(1 + \frac{t}{t_0}\right),$$

and so, for any $x \in [h(t) - \frac{\pi}{2}, h(t)]$, we have

$$Y(t) - x = \frac{3}{c_0} \ln\left(1 + \frac{t}{t_0}\right) + O(1) \in [0, \sqrt{t+1}\,], \quad \text{when } t \gg 1.$$

Using (47) we have

$$u(t, x) \leqslant Ct^{-\frac{5}{4}} \quad \text{when } t \gg 1, \tag{49}$$

where $C > 0$ are some constants depending on u_0, c_0, h_0. □

Remark 3.13. For any given $m \in (0, 1)$, denote

$$\chi(t) := \min\{x \in [g(t), h(t)] \mid u(t, x) = m\} \text{ and } \widetilde{\chi}(t) := \min\{y \in \mathbb{R} \mid \tilde{u}(t, y) = m/A\}.$$

Then by [21, Theorem 1.1] we have

$$c_0 t - \frac{3}{c_0} \ln t - C \leqslant \widetilde{\chi}(t) \leqslant c_0 t - \frac{3}{c_0} \ln t + C, \quad t \gg 1,$$

for some $C > 0$. Hence by (45) we have

$$\chi(t) \geqslant \beta t - \widetilde{\chi}(t) \geqslant (\beta - c_0)t + \frac{3}{c_0} \ln t - C, \quad t \gg 1. \tag{50}$$

4. Influence of β on the long time behavior of solutions

In this section we consider the influence of β on the long time behavior of the solutions. In Section 4.1 we give a locally uniformly convergence result. In Section 4.2 we consider the small advection $\beta \in (0, c_0)$ and prove Theorem 2.1. In Section 4.3 we first prove the boundedness of g_∞ for $\beta \geqslant c_0$, the boundedness of h_∞ for $\beta \geqslant \beta^*$, and then prove Theorem 2.4 for large advection $\beta \geqslant \beta^*$. In Section 4.4, we consider (P) with medium-sized advection $\beta \in [c_0, \beta^*)$ and prove Theorems 2.2 and 2.3. The arguments for the case $\beta \in [c_0, \beta^*)$ are longer and much more complicated than the cases with small or large advection.

4.1. Convergence result

First we give a locally uniformly convergence result for $\beta \geqslant c_0$.

Lemma 4.1. *Assume $\beta \geqslant c_0$. Then $u(t, \cdot)$ converges as $t \to \infty$ to 0 locally uniformly in I_∞.*

Proof. When $\beta > c_0$, the conclusion follows easily from (43) since the upper solution $Q_A(x - (\beta - c_0)t + x_0)$ is a rightward traveling wave with positive speed $\beta - c_0$ and $Q_A(z) \to 0$ as $z \to -\infty$.

Assume $\beta = c_0$. Then for any $[a, b] \subset I_\infty$, when t is sufficiently large we have

$$\frac{3}{c_0} \ln\left(1 + \frac{t}{t_0}\right) - \sqrt{t+1} < x < \frac{3}{c_0} \ln\left(1 + \frac{t}{t_0}\right) \quad \text{for any } x \in [a, b],$$

and so by (47), for any $x \in [a, b]$,

$$u(t, x) \leqslant C_4 e^{-\frac{5c_0}{12}\left[\frac{3}{c_0} \ln\left(1+\frac{t}{t_0}\right) - b\right]} \to 0, \quad \text{as } t \to \infty.$$

This proved the lemma. □

Theorem 4.2. *Let (u, g, h) be a time-global solution of (P). Then as $t \to \infty$, $u(t, \cdot)$ converges to 0 or to 1 locally uniformly in I_∞ when $\beta \in (0, c_0)$; $u(t, \cdot)$ converges to 0 locally uniformly in I_∞ when $\beta \geqslant c_0$.*

Moreover, $\lim_{t\to\infty} \|u(t, \cdot)\|_{L^\infty([g(t),h(t)])} = 0$ if I_∞ is a bounded interval.

Proof. When $\beta \geqslant c_0$, by Lemma 4.1 we have $u(t, \cdot) \to 0$ $(t \to \infty)$ locally uniformly in I_∞.

We now consider the case $\beta \in (0, c_0)$. Denote by $\omega(u)$ the ω-limit set of $u(t, \cdot)$ in the topology of $L^\infty_{\text{loc}}(I_\infty)$. In a similar way as proving Claim 1 in [13, Theorem 1.1] one can show that $\omega(u)$ consists of stationary solutions of $(P)_1$: 0, 1, $U(x - z_0)$, $U_l(x - z_0)$ and $W(x - z_0)$ for some $z_0 \in \mathbb{R}$ (see Section 3.2). If $U(x - z_0) \in \omega(u)$, then it is easy to show

by the monotonicity of $h(t)$ that $h_\infty = z_0$. Hence $U(x - z_0)$ is the unique element in $\omega(u)$ by the connectedness of $\omega(u)$. Moreover, by using the L^p estimates as in the proof of Claim 2 in [13, Theorem 1.1] one can show that $\|u(t,\cdot) - U(\cdot - z_0)\|_{C^{1+\nu}([0,h(t)])} \to 0$ as $t \to \infty$ for some $\nu \in (0,1)$. So

$$h'(t) = -\mu u_x(t, h(t)) \to -\mu U'(0) > 0 \quad \text{as } t \to \infty.$$

This contradicts $h(t) \to h_\infty = z_0$. This proves $U(x - z_0) \notin \omega(u)$. Similarly, one can show that $U_l(x - z_0), W(x - z_0) \notin \omega(u)$. Consequently, we have $\omega(u) = \{0\}$ or $\{1\}$.

Finally, when I_∞ is bounded, the uniform convergence $\|u(t,\cdot)\|_{L^\infty([g(t),h(t)])} \to 0$ can be proved in the same way as that in [13,28]. □

4.2. Problem with small advection: $0 < \beta < c_0$

In a similar way as proving [19, Lemma 2.2] and [13, Theorem 3.2, Corollary 4.5] one can show the following conditions for spreading and for vanishing.

Lemma 4.3. *Assume $\beta \in (0, c_0)$. Let (u, g, h) be a solution of (P).*

(i) *If $h_0 < H^* := \frac{\pi}{\sqrt{c_0^2 - \beta^2}}$ and if $\|u_0\|_{L^\infty([-h_0,h_0])}$ is sufficiently small, then vanishing happens;*
(ii) *if $h_0 \geqslant H^*$, then spreading happens.*

This lemma implies that, when $\beta \in (0, c_0)$, $2H^*$ is a critical width of the interval $I(t)$. Spreading happens if and only if $|I_\infty| > 2H^*$. This extends the results in [12,13] for $\beta = 0$, where it was shown that the critical width $2H^* = \frac{2\pi}{c_0} = \frac{\pi}{\sqrt{f'(0)}}$.

Proof of Theorem 2.1. By Lemma 4.3(ii) we see that spreading happens if $|I_\infty| > 2H^*$. By the definition of spreading, this implies that $I_\infty = \mathbb{R}$. Hence both the case $g_\infty > -\infty$, $h_\infty = \infty$ and the case $g_\infty = -\infty$, $h_\infty < \infty$ are impossible. I_∞ is either a bounded interval with width $|I_\infty| \leqslant 2H^*$ or the whole line $\mathbb{R}$. Using Theorem 4.2 and Lemma 4.3 again, we can get the spreading–vanishing dichotomy result for the long-time behavior of the solutions of (P). Then the sharp threshold of the initial data $\sigma\phi$ can be proved in a similar way as in [13, Theorem 5.2]. □

4.3. Boundedness of g_∞ and h_∞, the proof of Theorem 2.4

Whether g_∞ and h_∞ are bounded or not is also a part of the conclusions in the long time behavior of (u, g, h). In this subsection we show that $g_\infty > -\infty$ if $\beta \geqslant c_0$, and $h_\infty < \infty$ if $\beta \geqslant \beta^*$.

We will prove these conclusions by using Corollary 3.12. For this purpose we need the monotonicity of u_x. When $\beta = 0$, Du and Lou [13] proved the monotonicity of u_x in

$[h_0, h(t)]$ and in $[g(t), -h_0]$. When $\beta > 0$ we find that this is true only on the left side: $x \in [g(t), -h_0]$.

Lemma 4.4. *Assume (u, g, h) is a solution of (P). Then*

$$g(t) + h(t) > -2h_0 \quad \textit{for all } t > 0, \tag{51}$$

$$u_x(t, x) > 0 \quad \textit{for all } x \in [g(t), -h_0],\ t > 0. \tag{52}$$

Proof. It is easily seen by the continuity that, when $t > 0$ is sufficiently small, $g(t) + h(t) > -2h_0$ and $u_x(t, x) > 0$ for $g(t) \leqslant x \leqslant -h_0$. Define

$$T_1 := \sup\big\{s \mid g(t) + h(t) > -2h_0 \text{ for all } t \in (0, s)\big\},$$
$$T_2 := \sup\big\{s \mid u_x(t, x) > 0 \text{ for any } x \in [g(t), -h_0],\ t \in (0, s)\big\}.$$

We prove that $T_1 = T_2 = +\infty$. Otherwise, either $T_1 < T_2 \leqslant +\infty$, or $T_2 \leqslant T_1 \leqslant +\infty$ and $T_2 < +\infty$.

1. If $T_1 < T_2 \leqslant +\infty$, then

$$g(t) + h(t) > -2h_0 \text{ for } t \in [0, T_1) \quad \text{and} \quad g(T_1) + h(T_1) = -2h_0.$$

Hence

$$g'(T_1) + h'(T_1) \leqslant 0. \tag{53}$$

Set $G_{T_1} := \{(t, x) \mid t \in (0, T_1],\ x \in (g(t), -h_0)\}$ and

$$w(t, x) := u(t, x) - u(t, -2h_0 - x) \text{ in } \overline{G}_{T_1}.$$

Since $-h_0 \leqslant -2h_0 - x \leqslant -2h_0 - g(t) \leqslant h(t)$ when $(t, x) \in \overline{G}_{T_1}$, w is well-defined over $\overline{G}_{T_1}$ and it satisfies

$$w_t - w_{xx} - \beta w_x - c(t, x)w = -2\beta u_x(t, x) \leqslant 0 \text{ for } (t, x) \in G_{T_1},$$

where c is a bounded function, and

$$w(t, -h_0) = 0,\ w(t, g(t)) \leqslant 0 \text{ for } t \in (0, T_1].$$

Moreover,

$$w(T_1, g(T_1)) = u(T_1, g(T_1)) - u(T_1, -2h_0 - g(T_1)) = u(T_1, g(T_1)) - u(T_1, h(T_1)) = 0.$$

Then by the strong maximum principle and the Hopf lemma, we have

$$w(t, x) < 0 \text{ for } (t, x) \in G_{T_1},\ \text{and } w_x(T_1, g(T_1)) < 0.$$

Thus

$$g'(T_1) + h'(T_1) = -\mu[u_x(T_1, g(T_1)) + u_x(T_1, h(T_1))] = -\mu w_x(T_1, g(T_1)) > 0.$$

This contradicts (53).

2. If $T_2 \leqslant T_1 \leqslant \infty$ and $T_2 < +\infty$, then

$$u_x(t,x) > 0,\ t \in (0, T_2),\ x \in [g(t), -h_0].$$

By the definition of T_2, there exists $y \in (g(T_2), -h_0]$ such that $u_x(T_2, y) = 0$. Denote by x_0 the minimum of such y. By the continuity and the monotonicity of $g(t)$, there exists $T_0 \in [0, T_2)$ such that $x_0 = g(T_0)$. Let

$$G_{T_2} := \{(t,x) \mid t \in (T_0, T_2],\ x \in (g(t), x_0)\},$$
$$z(t,x) := u(t,x) - u(t, 2x_0 - x) \text{ for } (t,x) \in \overline{G}_{T_2}.$$

Using the maximum principle for $z(t,x)$ in G_{T_2} as above we conclude that $z_x(T_2, x_0) > 0$. This contradicts the definition of x_0.

Combining the above two steps we obtain $T_1 = T_2 = +\infty$. □

A direct consequence of Lemma 4.4 is $g_\infty + h_\infty \geqslant -2h_0$. So we have

Corollary 4.5. *There are only three possible situations for $I_\infty = (g_\infty, h_\infty)$:* (i) $I_\infty = \mathbb{R}$*;* (ii) I_∞ *is a finite interval; and* (iii) $I_\infty = (g_\infty, \infty)$ *with* $g_\infty > -\infty$.

Indeed, (i) and (ii) are possible when $\beta \in (0, c_0)$ (see Theorem 2.1), (ii) and (iii) are possible when $\beta \geqslant c_0$ (see Theorems 2.2, 2.3 and 2.4).

By the monotonicity of $u(t, \cdot)$ in $[g(t), -h_0]$, we can prove the boundedness of g_∞.

Proposition 4.6. *Assume $\beta \geqslant c_0$ and (u, g, h) is a solution of (P). Then $g_\infty > -\infty$.*

Proof. First we consider the case $\beta > c_0$. Let $f_A(s)$ be defined as in (42) and let $Q_A(z)$ be the unique solution of (4) with $c = c_0$, with f replaced by f_A and $q(+\infty) = 1$ replaced by $q(+\infty) = A$, as in (43). Since

$$Q_A(z) \sim -Cze^{\frac{c_0}{2}z} \text{ as } z \to -\infty \tag{54}$$

for some $C > 0$ (cf. [2,21]), there exist $T_1 > 0$, $C_1 > 0$ such that, when $t \geqslant T_1$,

$$\begin{aligned} Q_A(-h_0 - (\beta - c_0)t + x_0) &\leqslant -2C(-h_0 - (\beta - c_0)t + x_0)e^{\frac{c_0}{2}(-h_0 - (\beta - c_0)t + x_0)} \\ &\leqslant C_1 t e^{-\frac{c_0}{2}(\beta - c_0)t} \leqslant C_1 e^{-\frac{c_0}{4}(\beta - c_0)t}, \end{aligned}$$

where $x_0 > 0$ is large such that (43) holds. By Lemma 4.4 and (43) we have

$$u(t,x) \leqslant u(t,-h_0) \leqslant Q_A(-h_0-(\beta-c_0)t+x_0) \leqslant C_1 e^{-\frac{c_0}{4}(\beta-c_0)t} \tag{55}$$

for $x \in [g(t), -h_0]$ and $t \geqslant T_1$. Set $\delta := \min\{1, \frac{c_0}{4}(\beta-c_0)\}$, $\epsilon_1 := C_1 e^{-\frac{\beta\pi+c_0(\beta-c_0)T_1}{4}}$,

$$k(t) := -g(T_1) + \frac{\pi}{2} + \frac{\mu\epsilon_1}{\delta}(1-e^{-\delta t}) \text{ for } t \geqslant 0$$

and

$$w(t,x) := \epsilon_1 e^{-\delta t} e^{\frac{\beta}{2}(x+k(t))} \sin(x+k(t)) \quad \text{for } -k(t) \leqslant x \leqslant -k(t)+\frac{\pi}{2},\ t \geqslant 0.$$

A direct calculation shows that

$$\begin{aligned} w_t - w_{xx} + \beta w_x - f(w) &= w\Big[1-\delta+\frac{\beta}{2}k'(t)+\frac{\beta^2}{4}\Big] + k'(t)\hat{w} - f(w) \\ &\geqslant w\Big[1-\delta+\frac{\beta^2}{4}-f'(0)\Big] \geqslant (1-\delta)w \geqslant 0, \end{aligned}$$

for $-k(t) \leqslant x \leqslant -k(t)+\frac{\pi}{2}$, $t>0$, where $\hat{w} = \epsilon_1 e^{-\delta t} e^{\frac{\beta}{2}(x+k(t))}\cos(x+k(t))$,

$$-k'(t) = -\epsilon_1 \mu e^{-\delta t} = -\mu w_x(t,-k(t)), \quad t>0,$$

and

$$w\Big(t, -k(t)+\frac{\pi}{2}\Big) = \epsilon_1 e^{\frac{\beta\pi}{4}} e^{-\delta t} \geqslant C_1 e^{-\frac{c_0(\beta-c_0)}{4}(t+T_1)} \geqslant u(t+T_1, x)$$

for $g(t+T_1) \leqslant x \leqslant -h_0$, $t>0$. Hence for $t \geqslant 0$, either $g(t+T_1) \geqslant -k(t)+\frac{\pi}{2} \geqslant g(T_1) - \frac{\mu\epsilon_1}{\delta}$, or $u(t+T_1,\cdot)$ and $w(t,\cdot)$ have common domain. In the latter case, by comparing them on their common domain we have

$$g(t+T_1) \geqslant -k(t) \geqslant g(T_1) - \frac{\pi}{2} - \frac{\mu\epsilon_1}{\delta} > -\infty.$$

This proves $g_\infty > -\infty$.

Next we consider the case $\beta = c_0$. By Corollary 3.12 and Lemma 4.4, there exist $T_2 > \frac{5}{4}$ and $C>0$ such that

$$u(t,x) \leqslant u(t,-h_0) \leqslant Ct^{-\frac{5}{4}} \text{ for } g(t) \leqslant x \leqslant -h_0,\ t \geqslant T_2.$$

Set $\epsilon_2 := Ce^{-\frac{\beta\pi}{4}}$ and define

$$k_2(t) := -g(T_2) + \frac{\pi}{2} + 4\mu\epsilon_2[T_2^{-\frac{1}{4}} - (t+T_2)^{-\frac{1}{4}}] \text{ for } t>0,$$

$$w_2(t,x) := \epsilon_2(t+T_2)^{-\frac{5}{4}} e^{\frac{\beta}{2}(x+k_2(t))}\sin(x+k_2(t)) \text{ for } -k_2(t) \leqslant x \leqslant -k_2(t)+\frac{\pi}{2},\ t \geqslant 0.$$

A similar discussion as above shows that $(w_2, -k_2, -k_2 + \frac{\pi}{2})$ is an upper solution, and so

$$g(t + T_2) \geqslant -k_2(t) \geqslant g(T_2) - \frac{\pi}{2} - 4\mu\epsilon_2 T_2^{-\frac{1}{4}} > -\infty.$$

This proves the proposition. □

Next we prove the boundedness of h_∞ when $\beta \geqslant \beta^*$.

Proposition 4.7. *Assume $\beta \geqslant \beta^*$ and (u, g, h) is a solution of (P). Then $h_\infty < \infty$.*

Proof. 1. First we consider the case $\beta > \beta^*$. In this case we have $c^*(\beta) < \beta - c_0$. Denote $\nu := \beta - c_0 - c^*(\beta) > 0$. By (43), (41) and (54), there exist $T_1 > 0$, $C_1 > 0$ such that, for $x \in (g(t), h(t))$ and $t \geqslant T_1$, we have

$$\begin{aligned} u(t,x) &\leqslant Q_A(x - (\beta - c_0)t + x_0) \leqslant Q_A(h(t) - (\beta - c_0)t + x_0) \\ &\leqslant Q_A(-\nu t + H + x_0) \leqslant -2C(-\nu t + H + x_0)e^{\frac{c_0}{2}(-\nu t + H + x_0)} \\ &\leqslant C_1 e^{-\frac{c_0\nu}{4}t}. \end{aligned}$$

Set $\delta := \frac{1}{2}\min\{1, \frac{c_0\nu}{4}\}$ and choose $T_2 > T_1$ such that

$$\epsilon_3 := C_1 e^{\frac{\beta\pi - c_0\nu T_2}{4}} < \frac{1}{\beta\mu}.$$

Define

$$k_3(t) := h(T_2) + \frac{\pi}{2} + \frac{\mu\epsilon_3}{\delta}(1 - e^{-\delta t}) \text{ for } t \geqslant 0$$

and

$$w_3(t,x) := \epsilon_3 e^{-\delta t} e^{\frac{\beta}{2}(x - k_3(t))} \cos\left(x - k_3(t) + \frac{\pi}{2}\right) \text{ for } k_3(t) - \frac{\pi}{2} \leqslant x \leqslant k_3(t),\ t \geqslant 0.$$

A direct calculation as in the proof of Proposition 4.6 shows that $(w_3, k_3(t) - \frac{\pi}{2}, k_3(t))$ is an upper solution and

$$h(t + T_2) \leqslant k_3(t) \leqslant h(T_2) + \frac{\pi}{2} + \frac{\mu\epsilon_3}{\delta} < \infty.$$

2. Next we consider the case $\beta = \beta^*$. We first show that, for some large T_3,

$$u(t,x) < U^*(x - c^*(\beta^*)t + h_0) \text{ for } x \in [g(t), h(t)],\ t \geqslant T_3, \tag{56}$$

where $U^*(x - c^*(\beta^*)t + h_0)$ is the rightward traveling semi-wave with endpoint at $c^*(\beta^*)t - h_0$. At time $t = 0$, $u(t,x)$ and $U^*(x + h_0)$ intersect at $x = -h_0$. Then for small time $t > 0$, they intersect at exactly one point.

We claim that the case $c^*(\beta^*)t - h_0 < h(t)$ for all $t \geqslant 0$ is impossible. Otherwise, combining with (41) we have $h(t) = c^*(\beta^*)t + O(1)$, and so by Corollary 3.12 there exist $T_4 > 0$ and $C > 0$ such that

$$u(t,x) \leqslant Ct^{-\frac{5}{4}} \text{ for } h(t) - \frac{\pi}{2} \leqslant x \leqslant h(t),\ t \geqslant T_4.$$

Set $\epsilon_4 := Ce^{\frac{\beta\pi}{4}}$, $T_5 := \max\{T_4, \frac{5}{4} + \frac{\beta\mu\epsilon_4}{2}\}$ and define

$$k_4(t) := h(T_5) + \frac{\pi}{2} + 4\mu\epsilon_4[T_5^{-\frac{1}{4}} - (t+T_5)^{-\frac{1}{4}}], \quad t \geqslant 0,$$

$$w_4(t,x) := \epsilon_4(t+T_5)^{-\frac{5}{4}} e^{\frac{\beta}{2}(x-k_4(t))} \cos\left(x - k_4(t) + \frac{\pi}{2}\right), \quad k_4(t) - \frac{\pi}{2} \leqslant x \leqslant k_4(t),\ t \geqslant 0.$$

A direct calculation shows that $(w_4, k_4(t) - \frac{\pi}{2}, k_4(t))$ is an upper solution, and so

$$h(t+T_5) \leqslant k_4(t) \leqslant h(T_5) + \frac{\pi}{2} + 4\mu\epsilon_4 T_5^{-\frac{1}{4}} < \infty,$$

contradicts our assumption $c^*(\beta^*)t - h_0 < h(t)$ for all t.

Therefore, there exists $T_6 > 0$ such that $h(T_6) = c^*(\beta^*)T_6 - h_0$ and by Lemma 3.10, the unique intersection point between u and U^* disappears after T_6, and so (56) holds for any $T_3 > T_6$. (This can also be proved by the fact $u(T_6, x) \leqslant U^*(x - c^*(\beta^*)T_6 + h_0)$ for $x \in [g(T_6), h(T_6)]$ and by the comparison principle.)

On the other hand, by Lemma 3.6, $V^*_{\delta_1}(z) := V(z; \beta^* - c_0 - \delta_1, -c_0 - \delta_1)$ approaches $U^*(z)$ locally uniformly in $(-\infty, 0]$ as $\delta_1 \to 0$. Hence there exists $\delta_1 > 0$ sufficiently small such that $V^*_{\delta_1}(z)$ is close to $U^*(z)$ and so

$$u(T_3, x) < V^*_{\delta_1}(x - c^*(\beta^*)T_3 + h_0) \text{ for } x \in [g(t), h(t)].$$

By comparison $u(t+T_3, x) \leqslant V^*_{\delta_1}(x - (\beta^* - c_0 - \delta_1)t - c^*(\beta^*)T_3 + h_0)$ and so $h(t+T_3)$ is blocked by the right endpoint $(\beta^* - c_0 - \delta_1)t + c^*(\beta^*)T_3 - h_0$ of $V^*_{\delta_1}$:

$$h(t+T_3) \leqslant (\beta^* - c_0 - \delta_1)t + c^*(\beta^*)T_3 - h_0, \quad t \geqslant 0.$$

Using (43) we see that, for any $x \in [g(t), h(t)]$ and sufficiently large t,

$$u(t,x) \leqslant Q_A(x - (\beta^* - c_0)t + x_0) \leqslant Q_A(-\delta_1 t + x_1) \leqslant C_2 e^{-\frac{c_0\delta_1}{4}t},$$

for some $x_1 \in \mathbb{R}$ and $C_2 > 0$. The rest proof is similar as that in step 1.

This proves the proposition. □

Proof of Theorem 2.4. The conclusions follow from Propositions 4.6, 4.7 and Theorem 4.2 immediately. □

4.4. Problem with medium-sized advection: $c_0 \leqslant \beta < \beta^*$

In this subsection we consider the case $\beta \in [c_0, \beta^*)$. In this case, the long time behavior of the solutions is complicated and more interesting. Besides vanishing, we find some new phenomena: virtual spreading, virtual vanishing and convergence to the tadpole-like traveling wave.

In the first part, we give some sufficient conditions for vanishing; in the second part we give a necessary and sufficient condition for virtual spreading; in the third part we study the limits of $h'(t)$ and $u(t,x)$ when vanishing and virtual spreading do not happen; in the last part we finish the proofs of Theorems 2.2 and 2.3.

4.4.1. Vanishing phenomena

When $\beta \geqslant c_0$, we have $g_\infty > -\infty$ by Proposition 4.6, which implies that $u \to 0$ locally uniformly. We now show that the convergence can be a uniform one when the initial data u_0 is sufficiently small.

Lemma 4.8. *Assume $\beta \geqslant c_0$ and (u,g,h) is the solution of (P). If $\|u_0\|_{L^\infty([-h_0,h_0])}$ is sufficiently small, then vanishing happens.*

Proof. Choose $\delta > 0$ small such that

$$\frac{\pi^2}{h_0^2(1+\delta)^2} \geqslant 4\delta + \beta h_0 \delta^2.$$

Set $k(t) := h_0(1+\delta-\frac{\delta}{2}e^{-\delta t})$, $\epsilon := \frac{h_0^2\delta^2}{\pi\mu}(1+\frac{\delta}{2})$ and

$$w(t,x) := \epsilon e^{-\delta t} e^{\frac{\beta}{2}(x-k(t))} \cos\frac{\pi x}{2k(t)} \quad \text{for } -k(t) \leqslant x \leqslant k(t),\ t>0. \tag{57}$$

A direct calculation shows that, for $x \in (-k(t), k(t))$ and $t > 0$,

$$w_t - w_{xx} + \beta w_x - f(w) \geqslant \frac{1}{4}\Big(\frac{\pi^2}{h_0^2(1+\delta)^2} - 4\delta - \beta h_0\delta^2\Big) w \geqslant 0.$$

On the other hand, for any $t > 0$,

$$\mu w_x(t,-k(t)) \leqslant -\mu w_x(t,k(t)) = \frac{\pi\mu\epsilon}{2k(t)} e^{-\delta t} \leqslant \frac{\pi\mu\epsilon}{2h_0(1+\frac{\delta}{2})} e^{-\delta t} = \frac{\delta^2}{2} h_0 e^{-\delta t} = k'(t).$$

Hence $(w,-k,k)$ is an upper solution of (P). Clearly $k(0) = h_0(1+\frac{\delta}{2}) > h_0$ and $w(t,\pm k(t)) = 0$ for $t \geqslant 0$. If $\|u_0\|_{L^\infty([-h_0,h_0])}$ is small such that

$$\|u_0\|_{L^\infty([-h_0,h_0])} \leqslant \epsilon e^{-\frac{\beta}{2}h_0(2+\frac{\delta}{2})} \cos\frac{\pi}{2+\delta} = w(0,-h_0),$$

then $u_0(x) \leqslant w(0,x)$ for $x \in [-h_0, h_0]$. By the comparison principle, we have

$$-h_0(1+\delta) \leqslant -k(t) \leqslant g(t) < h(t) \leqslant k(t) \leqslant h_0(1+\delta),$$
$$\|u(t,\cdot)\|_{L^\infty([g(t),h(t)])} \leqslant \|w(t,\cdot)\|_{L^\infty([-k(t),k(t)])} \leqslant \epsilon e^{-\delta t} \to 0 \quad \text{as } t \to \infty.$$

This proves the lemma. □

For any given $h_0 > 0$ and $\phi \in \mathscr{X}(h_0)$, we write the solution (u, g, h) also as $(u(t,x;\sigma\phi), g(t;\sigma\phi), h(t;\sigma\phi))$ to emphasize the dependence on the initial data $u_0 = \sigma\phi$. Set

$$E_0 := \{\sigma \geqslant 0 \mid \text{vanishing happens for } u(\cdot,\cdot;\sigma\phi)\}, \quad \sigma_* := \sup E_0. \tag{58}$$

Lemma 4.8 implies that $\sigma \in E_0$ for all small $\sigma > 0$. By the comparison principle we have $[0, \sigma_*) \subset E_0$. In case $\sigma_* = \infty$ (this happens in particular when $\liminf_{s\to\infty} \frac{-f(s)}{s^8} = \infty$ and $\beta = 0$, see [13, Proposition 5.4]), there is nothing left to prove. Hence we only consider the case $\sigma_* \in (0, \infty)$.

Theorem 4.9. *Assume $c_0 \leqslant \beta < \beta^*$. For any $\phi \in \mathscr{X}(h_0)$, let E_0 and σ_* be defined as in (58). If $\sigma_* \in (0,\infty)$, then $E_0 = [0, \sigma_*)$. If $\sigma \geqslant \sigma_*$, then $g(\infty;\sigma\phi) > -\infty$, $h(\infty;\sigma\phi) = \infty$, and $u(t,\cdot;\sigma\phi) \to 0$ locally uniformly in $(g(\infty;\sigma\phi), \infty)$.*

Proof. For any positive $\sigma_0 \in E_0$, since $u(t,\cdot;\sigma_0\phi) \to 0$ uniformly, we can find a large $T_0 > 0$ such that $u(T_0, x; \sigma_0\phi) < w(0,x)$, where $w(t,x)$ is defined as in (57), with h_0 replaced by $H := \max\{h(\infty;\sigma_0\phi), -g(\infty;\sigma_0\phi)\} < \infty$. By continuity, there exists $\epsilon > 0$ such that $u(T_0, x; \sigma\phi) < w(0,x)$ for every $\sigma \in [\sigma_0, \sigma_0 + \epsilon)$. As in the proof of Lemma 4.8, we conclude that vanishing happens for $u(t,x;\sigma\phi)$, that is, $\sigma \in E_0$. Therefore, $E_0 \backslash \{0\}$ is an open set, and so $E_0 = [0, \sigma_*)$. The rest of the conclusions follow from Theorem 4.2 and Proposition 4.6. □

We finish this part by giving another sufficient condition for vanishing.

Lemma 4.10. *Assume $c_0 < \beta < \beta^*$. Let (u, g, h) be a solution of (P). Then vanishing happens if there exist $t_1 \geqslant 0$, $x_1 \in \mathbb{R}$ such that*

$$h(t_1) \leqslant x_1, \ u(t_1, x) \leqslant V^*(x - x_1) \text{ for } x \in [g(t_1), h(t_1)],$$

where $V^*(z)$ is the tadpole-like solution as in Lemma 3.5(ii).

Proof. Since $V^*(x - (\beta - c_0)t - x_1)$ is a solution of $(P)_1$ satisfying Stefan free boundary condition at $x = k(t) := (\beta - c_0)t + x_1$:

$$k'(t) = \beta - c_0 = -\mu(V^*)'(0)$$

by Lemma 3.5(ii). By the comparison principle we have

$$h(t_1+1) < k(1),\ u(t_1+1,x) < V^*(x-\beta+c_0-x_1) \text{ for } x \in [g(t_1+1), h(t_1+1)].$$

By Lemma 3.5(iii), $V_\delta(z) := V(z;\beta-c_0-\delta,-c_0-\delta) \to V^*(z)$ locally uniformly in $(-\infty,0]$ as $\delta \to 0$. Hence for sufficiently small $\delta > 0$, we have

$$u(t_1+1,x) < V_\delta(x-\beta+c_0-x_1) \text{ for } x \in [g(t_1+1), h(t_1+1)].$$

Since $V_\delta(x-(\beta-c_0-\delta)t-\beta+c_0-x_1)$ is a solution of $(P)_1$ satisfying Stefan free boundary condition at $x = k_\delta(t) := (\beta-c_0-\delta)t+\beta-c_0+x_1$, by comparison we have

$$h(t+t_1+1) \leqslant k_\delta(t) = (\beta-c_0-\delta)t+\beta-c_0+x_1.$$

A similar argument as in step 1 of the proof of Proposition 4.7 shows that $h_\infty < \infty$, this implies that vanishing happens by Theorem 4.2. □

4.4.2. A necessary and sufficient condition for virtual spreading

Lemma 4.11. *Assume $c_0 \leqslant \beta < \beta^*$. Let (u,g,h) be a solution of (P). Then virtual spreading happens if and only if, for any $\delta \in (0, c^*(\beta)-\beta+c_0)$, there exist t_1 and x_1 such that*

$$u(t_1,x) \geqslant W_\delta(x-x_1) \text{ for } x \in [x_1-L_\delta, x_1], \tag{59}$$

where $W_\delta(z)$, $L_\delta := L(\beta-c_0+\delta,-c_0+\delta)$ are defined in Lemma 3.7.

Proof. The inequality (59) follows from the definition of u being virtual spreading immediately. We only need to show that (59) is a sufficient condition for virtual spreading.

Since $W_\delta(x-(\beta-c_0+\delta)t-x_1)$ satisfies $(P)_1$ and Stefan free boundary condition at $x = r(t) := (\beta-c_0+\delta)t+x_1$, comparing u and W_δ we have

$$u(t+t_1,x) > W_\delta(x-(\beta-c_0+\delta)t-x_1) \text{ for } x \in [r(t)-L_\delta, r(t)],\ t > 0. \tag{60}$$

In particular, this is true at $t = 1$. Since $W_\delta(z)$ depends on δ continuously, we have

$$u(t_1+1,x) > W_{\delta+\epsilon}(x-(\beta-c_0+\delta+\epsilon)-x_1) \text{ for } x \in [r(1)+\epsilon-L_{\delta+\epsilon}, r(1)+\epsilon],$$

for any $\epsilon \in (0,\epsilon_0]$ provided $\epsilon_0 > 0$ is small. Using the comparison principle again between $u(t+t_1+1,x)$ and $W_{\delta+\epsilon}(x-(\beta-c_0+\delta+\epsilon)(t+1)-x_1)$, we have

$$h(t+t_1+1) \geqslant (\beta-c_0+\delta+\epsilon)(t+1)+x_1,\ t > 0.$$

This implies that

$$H(t) := h(t+t_1+1) - (\beta - c_0 + \delta)t \geqslant \epsilon(t+1) + x_1 \to \infty \text{ as } t \to \infty. \tag{61}$$

Set

$$G(t) := g(t+t_1+1) - (\beta - c_0 + \delta)t \tag{62}$$

and

$$w(t,x) := u(t+t_1+1, x + (\beta - c_0 + \delta)t) \text{ for } G(t) \leqslant x \leqslant H(t),\ t \geqslant 0. \tag{63}$$

Then $G(t) \to -\infty$ as $t \to \infty$ by Proposition 4.6, w satisfies

$$\begin{aligned} &w(t,x) > W_\delta(x - (\beta - c_0 + \delta) - x_1) \\ &\quad \text{for } x \in [x_1 + \beta - c_0 + \delta - L_\delta, x_1 + \beta - c_0 + \delta],\ t \geqslant 0 \end{aligned} \tag{64}$$

by (60) and

$$\begin{cases} w_t = w_{xx} - (c_0 - \delta)w_x + f(w), \quad t > 0,\ G(t) < x < H(t), \\ w(t, G(t)) = 0, \quad G'(t) = -\mu w_x(t, G(t)) - (\beta - c_0 + \delta), \quad t > 0, \\ w(t, H(t)) = 0, \quad H'(t) = -\mu w_x(t, H(t)) - (\beta - c_0 + \delta), \quad t > 0, \\ G(0) = g(t_1+1),\ H(0) = h(t_1+1), \quad w(0,x) = u(t_1+1, x) \text{ for } G(0) \leqslant x \leqslant H(0). \end{cases} \tag{65}$$

Since $H(t), -G(t) \to \infty$ as $t \to \infty$, for any large $L > 0$, there exists $T_L > 0$ such that $H(t), -G(t) > L$ for all $t \geqslant T_L$. Let $\tilde{w}_L(t,x)$ be the solution of

$$\begin{cases} \tilde{w}_t = \tilde{w}_{xx} - (c_0 - \delta)\tilde{w}_x + f(\tilde{w}), & x \in (-L, L),\ t > 0, \\ \tilde{w}(t, \pm L) = 0, & t > 0, \\ \tilde{w}(0,x) = w(T_L, x), & x \in [-L, L]. \end{cases}$$

Then, by the comparison principle we have $w(t+T_L, x) \geqslant \tilde{w}_L(t,x)$ for $x \in [-L, L]$. Since f is a Fisher-KPP type of nonlinearity, it is well known that $\tilde{w}_L(t,\cdot)$ converges as $t \to \infty$ to the unique positive solution $\bar{w}_L(x)$ of

$$\bar{w}_{xx} - (c_0 - \delta)\bar{w}_x + f(\bar{w}) = 0 \ \ (x \in (-L, L)), \qquad \bar{w}(\pm L) = 0.$$

Hence $\liminf_{t\to\infty} w(t+T_L, x) \geqslant \bar{w}_L(x)$ for $x \in [-L, L]$. Since this is true for any large L and since $\bar{w}_L(x) \to 1$ as $L \to \infty$ locally uniformly in $x \in \mathbb{R}$, we deduce that $\liminf_{t\to\infty} w(t,x) \geqslant 1$ for all $x \in \mathbb{R}$. On the other hand, by the property (F) we have $\limsup_{t\to\infty} w(t,x) \leqslant 1$, and so $w(t,\cdot) \to 1$ as $t \to \infty$ locally uniformly in $x \in \mathbb{R}$. This means that spreading happens for w and so virtual spreading happens for u. This proves the lemma. $\square$

4.4.3. The limits of h and u when vanishing and virtual spreading do not happen

In this part we always assume $c_0 \leqslant \beta < \beta^*$ and $\sigma_* \in (0, \infty)$ for given $\phi \in \mathscr{X}(h_0)$, where σ_* is defined by (58). We consider the limits of $h(t; \sigma\phi)$, $h'(t; \sigma\phi)$ and $u(t, \cdot + h(t); \sigma\phi)$ when vanishing does not happen, that is, when $\sigma \geqslant \sigma_*$.

Lemma 4.12. *Assume $c_0 \leqslant \beta < \beta^*$. If vanishing does not happen for a solution u of (P), then*

$$\lim_{t\to\infty} [h(t) - (\beta - c_0)t] = +\infty. \tag{66}$$

Proof. When $\beta = c_0$, we have $g_\infty > -\infty$ by Proposition 4.6. If $h_\infty < \infty$, then vanishing happens for u by Theorem 4.2, contradicting our assumption. Therefore, (66) holds when $\beta = c_0$.

We now consider the case $c_0 < \beta < \beta^*$. First we prove that

$$h(t) > (\beta - c_0)t - h_0 \text{ for any } t > 0.$$

Set

$$\eta_1(t, x) := u(t, x) - V^*(x - (\beta - c_0)t + h_0) \text{ for } x \in J_1(t),\ t > 0,$$

where

$$J_1(t) := [g(t), \min\{h(t), (\beta - c_0)t - h_0\}] \text{ for } t > 0.$$

It is easily seen that, for $0 < t \ll 1$,

$$(\beta - c_0)t - h_0 < h(t) \text{ and } \mathcal{Z}_{J_1(t)}[\eta_1(t, \cdot)] = 1. \tag{67}$$

We claim that this is true for all $t > 0$. Otherwise, there exists $T_1 > 0$ such that $(\beta - c_0)t - h_0 < h(t)$ for $0 < t < T_1$ and $(\beta - c_0)T_1 - h_0 = h(T_1)$. By Lemma 3.10 we have $\mathcal{Z}_{J_1(t)}[\eta_1(t, \cdot)] = 1$ for $0 < t < T_1$ and $\mathcal{Z}_{J_1(t)}[\eta_1(t, \cdot)] = 0$ for $t > T_1$. Therefore,

$$u(t, x) < V^*(x - (\beta - c_0)t + h_0) \text{ for } x \in I(t),\ T_1 < t \ll T_1 + 1.$$

This implies that vanishing happens for u by Lemma 4.10, contradicting our assumption.

Next we prove that, for any large $M > 0$, $h(t) > (\beta - c_0)t + M$ when t is large. By the definition of $\mathscr{X}(h_0)$, there exists $X > h_0$ large such that $u_0(x)$ intersects $V^*(x - M)$ at exactly two points for any $M \geqslant X$. Set

$$\eta_2(t, x) := u(t, x) - V^*(x - (\beta - c_0)t - M) \text{ for } x \in J_2(t),\ t > 0,$$

where

$$J_2(t) := [g(t), \min\{h(t), (\beta - c_0)t + M\}].$$

Then $\mathcal{Z}_{J_2(t)}[\eta_2(t,\cdot)] = 2$ for $0 < t \ll 1$. Denote by $\xi_1(t)$ and $\xi_2(t)$ with $\xi_1(t) < \xi_2(t)$ the two zeros of $\eta_2(t,\cdot)$. Then we have the following situations about the relations among $\xi_1(t)$, $\xi_2(t)$, $h(t)$ and $(\beta - c_0)t + M$.

Case 1. $h(t) < (\beta - c_0)t + M$ for all $t > 0$. In this case, combining with (67) we have $h(t) = (\beta - c_0)t + O(1)$. Using a similar argument as in step 2 of the proof of Proposition 4.7 we can derive $h_\infty < \infty$. This implies that vanishing happens for u, contradicting our assumption.

Case 2. There exists $T_2 > 0$ such that $h(t) < (\beta - c_0)t + M$ for $0 < t < T_2$ and $h(T_2) = (\beta - c_0)T_2 + M$. This includes several subcases.

Subcase 2-1. $\xi_1(t)$ meets $\xi_2(t)$ at time $t = T_3 < T_2$. In this case, $\xi_1(T_3) = \xi_2(T_3)$ is a degenerate zero of $\eta_2(T_3,\cdot)$ and so $\mathcal{Z}_{I(t)}[\eta_2(t,\cdot)] = 0$ for $T_3 < t \ll T_3 + 1$. This indicates that

$$u(t,x) < V^*(x - (\beta - c_0)t - M), \quad x \in [g(t), h(t)],\ T_3 < t \ll T_3 + 1, \tag{68}$$

and so vanishing happens by Lemma 4.10, contradicting our assumption.

Subcase 2-2. $\xi_1(t) < \xi_2(t) < h(t)$ for $0 < t \leqslant T_2$. This means a new intersection point $(h(T_2), 0)$ between u and V^* emerges on the boundary. This is impossible by Lemma 3.10.

Subcase 2-3. $\xi_1(t) < \xi_2(t) < h(t)$ for $0 < t < T_2$ and $\xi_1(T_2) = \xi_2(T_2) = h(T_2)$. This means the two intersection points between u and V^* move rightward to $(h(t), 0)$ at time T_2. By Lemma 3.10, this is the unique zero of $\eta_2(T_2,\cdot)$ and it will disappear after time T_2. Hence (68) holds for $t > T_2$. Then vanishing happens, a contradiction.

Subcase 2-4. $\xi_1(t) < \xi_2(t) < h(t)$ for $0 < t < T_2$ and $\xi_1(T_2) < \xi_2(T_2) = h(T_2) = (\beta - c_0)T_2 + M$. By Lemma 3.10, $\mathcal{Z}_{J_2(t)}[\eta_2(t,\cdot)] = 1 < 2$ for $T_2 < t \ll T_2 + 1$, where $J_2(t) := [g(t), (\beta - c_0)t + M]$. We claim that

$$(\beta - c_0)t + M < h(t) \text{ for all } t > T_2 \tag{69}$$

and so $\mathcal{Z}_{J_2(t)}[\eta_2(t,\cdot)] = 1$ for all $t > T_2$. Indeed, if $(\beta - c_0)t + M$ catches up $h(t)$ again at time $t = T_4 > T_2$, then $h(T_4)$ is a zero of η_2, by Lemma 3.10 it cannot be a new one and must be ξ_1, that is, the unique intersection point $(\xi_1(t), u(t, \xi_1(t)))$ (for $t \in [T_2, T_4)$) moves to $(h(t), 0)$ at time T_4. Moreover, this unique zero will disappear after T_4 by Lemma 3.10. This implies that (68) holds for $t > T_4$ and so vanishing happens, a contradiction. Consequently, (69) holds for any $M \geqslant X$ and so (66) holds. □

Lemma 4.13. *Assume $c_0 \leqslant \beta < \beta^*$. If vanishing and virtual spreading do not happen for the solution u of (P), then*

$$\lim_{t\to\infty} h'(t) = \beta - c_0. \tag{70}$$

Proof. We divide the proof into several steps.

Step 1. We first prove $h'(t) \geqslant \beta - c_0$ for all large t. This is clear when $\beta = c_0$. We now assume $c_0 < \beta < \beta^*$.

For readers' convenience, we first sketch the idea of our proof. We put a tadpole-like traveling wave $V^*(x - (\beta - c_0)t - C)$ whose right endpoint $r(t) := (\beta - c_0)t + C$ lies right to h_0. As t increasing, both $h(t)$ and $r(t)$ move rightward, but $h(t)$ moves faster by Lemma 4.12. Hence $h(t)$ catches up $r(t)$ at some time T. We will show that at this moment $u > V^*$ near $x = h(T)$ and so $h'(T) \geqslant \beta - c_0$. Since the shift C of V^* can be chosen continuously we indeed obtain $h'(t) \geqslant \beta - c_0$ for all large time t.

Now we give the details of the proof. As in the proof of the previous lemma, there exists $X > h_0$ such that $u_0(x)$ intersects $V^*(x-M)$ at exactly two points for any $M \geqslant X$.

By (66), there exists $T_X > 0$ such that $h(t) - (\beta - c_0)t > X$ for all $t \geqslant T_X$. For any $a > h(T_X)$ denote by T_a the unique time such that $h(T_a) = a$. Set $X_a := h(T_a) - (\beta - c_0)T_a\ (> X)$. We study the intersection points between $u(t, \cdot)$ and $V^*(x-(\beta-c_0)t-X_a)$. As in the proof of the previous lemma, only Subcase 2-4 is possible, that is, there exists $T^* > 0$ such that

$$\begin{cases} \text{when } 0 < t < T^*, \quad \xi_1(t) < \xi_2(t) < h(t) < (\beta - c_0)t + X_a, \\ \text{when } t = T^*, \quad \xi_1(t) < \xi_2(t) = h(t) = (\beta - c_0)t + X_a, \\ \text{when } t > T^*, \quad h(t) > (\beta - c_0)t + X_a. \end{cases}$$

Therefore T^* is nothing but T_a. At this moment we have

$$u(T_a, x) > V^*(x - (\beta - c_0)T_a - X_a) \text{ for } x \in (\xi_1(T_a), h(T_a)),$$

and

$$h'(T_a) = -\mu u_x(T_a, h(T_a)) \geqslant -\mu(V^*)'(0) = \beta - c_0.$$

Since $a > h(T_X)$ is arbitrary, T_a is continuous and strictly increasing in a, we indeed have $h'(t) \geqslant \beta - c_0$ for all $t > T_X$.

Step 2. We prove

$$\lim_{t\to\infty}[h(t) - (\beta - c_0 + \delta)t] = -\infty \text{ for all } \delta \in (0, c^* - \beta + c_0). \tag{71}$$

For any $\delta \in (0, c^* - \beta + c_0)$, we choose $\delta_1 \in (0, \delta)$ and consider the compactly supported traveling wave $W_{\delta_1}(x - c_1 t - M)$, where $c_1 = \beta - c_0 + \delta_1$, $M > 0$ is a large real number such that $u_0(x)$ has no intersection point with $W_{\delta_1}(x - M)$. Clearly (71) is proved if we have $h(t) < c_1 t + M$ for all $t > 0$. If, otherwise, there exists some $T_1 > 0$ such that

$$h(t) < c_1 t + M \text{ for } t \in [0, T_1), \ h(T_1) = c_1 T_1 + M,$$

then there exists $T_2 \in (0, T_1)$ such that $h(t)$ catches up the left boundary $l_1(t) := c_1 t + M - L_{\delta_1}$ of the support of $W_\delta(x - c_1 t - M)$ at time T_2 and never lags behind it again. So in the time interval (T_2, T_1).

$$\mathcal{Z}_{J_1(t)}[\zeta_1(t, \cdot)] = 1 \text{ for } t \in (T_2, T_1),$$

where $J_1(t) := [l_1(t), h(t)]$ and

$$\zeta_1(t, x) := u(t, x) - W_{\delta_1}(x - c_1 t - M) \text{ for } x \in J_1(t),\ t \in [T_2, T_1].$$

By Lemma 3.10, the unique zero $\zeta_1(t, \cdot)$ tends to $h(T_1)$ as $t \to T_1$ and it disappears after T_1. Hence

$$u(T_1, x) \geqslant W_{\delta_1}(x - c_1 T_1 - M) \text{ for } x \in [l_1(T_1), c_1 T_1 + M] = [l(T_1), h(T_1)].$$

This implies that virtual spreading happens for u by Lemma 4.11, contradicts our assumption.

Step 3. Based on Step 2 we prove

$$h'(t) \leqslant \beta - c_0 + \delta \text{ for large } t, \tag{72}$$

for any $\delta \in (0, c^* - \beta + c_0)$. Fix such a δ, we consider $u(t, x)$ and $W_\delta(x - (\beta - c_0 + \delta)t + h_0)$. Without loss of generality, we assume that $h(0) - g(0) = 2h_0$ is bigger than the width L_δ of the support of W_δ, for otherwise, we can replace $u(t, x)$ by $u(t + T, x)$ for some large T to carry out the following argument.

It is easily seen that these two functions intersect at exactly one point in their common domain $J_2(t) := [g(t), r(t)]$ for small $t > 0$, where $r(t) := (\beta - c_0 + \delta)t - h_0$. By Step 2, there exists $T_3 > 0$ such that

$$r(t) < h(t) \text{ for } t \in [0, T_3),\ r(T_3) = h(T_3).$$

Denote the left boundary of the support of $W_\delta(x - r(t))$ by $l_2(t) := r(t) - L_\delta$. Then $l_2(T_3) = h(T_3) - L_\delta > h(0) - L_\delta > g(0) > g(T_3)$. Hence $l_2(t)$ meets $g(t)$ before T_3, that is, there exists $T_4 \in (0, T_3)$ such that

$$l_2(t) < g(t) \text{ for } t \in [0, T_4),\ l_2(T_4) = g(T_4).$$

It follows from Lemma 3.10 that, when $T_4 < t \ll T_4 + 1$, either $W_\delta(x - r(t)) < u(t, x)$ in $[l_2(t), r(t)]$ or $\mathcal{Z}_{[l_2(t), r(t)]}[u(t, \cdot) - W_\delta(\cdot - r(t))] = 2$ holds. In the former case, virtual spreading happens for u by Lemma 4.11, contradicting our assumption. In the latter case, a similar discussion as in the proof of the previous lemma shows that the only possibility is that $r(t)$ catches up $h(t)$ at $t = T_3$, and the other intersection point $\xi(T_3)$ between $u(T_3, \cdot)$ and $W_\delta(\cdot - r(T_3))$ stays on the left. Hence we have

$$u(T_3, x) < W_\delta(x - r(T_3)) \text{ for } x \in (\xi(T_3), h(T_3)),$$

and so

$$h'(T_3) = -\mu u_x(T_3, h(T_3)) \leqslant -\mu W'_\delta(0) = \beta - c_0 + \delta. \tag{73}$$

Using the similar idea as in Step 1 of the current proof, we obtain (72) for all large time t.

Step 4. Combining Step 1 with Step 3 we have

$$\beta - c_0 \leqslant h'(t) \leqslant \beta - c_0 + \delta \text{ for large } t.$$

Since $\delta > 0$ can be arbitrarily small, we proves (70). □

Lemma 4.14. *Under the assumption of Lemma 4.13, $u(t, \cdot)$ has exactly one local maximum point for large t.*

Proof. Using zero number argument Lemma 3.9 to $u_x(t, \cdot)$ we see that $u(t, \cdot)$ has exactly N local maximum points for large t, where N is a positive integer. If $N \geqslant 2$, then by Lemma 3.11 the leftmost maximum point $\xi_1(t)$ moves right at a speed not less than β. On the other hand, (70) indicates $h(t)$ moves right at a speed $\beta - c_0$. Therefore, after some time, $\xi_1(t)$ reaches $h(t)$, this is a contradiction. □

Theorem 4.15. *Assume that vanishing and virtual spreading do not happen for the solution u of (P).*

(i) *If $c_0 < \beta < \beta^*$, then*

$$\lim_{t\to\infty} \|u(t, \cdot) - V^*(\cdot - h(t))\|_{L^\infty(I(t))} = 0; \tag{74}$$

(ii) *If $\beta = c_0$, then*

$$\lim_{t\to\infty} \|u(t, \cdot)\|_{L^\infty(I(t))} = 0; \tag{75}$$

Proof. 1. We first prove the locally uniform convergence near $h(t)$. Set $w(t, x) := u(t, x + h(t))$ and $G(t) := g(t) - h(t)$ for $t \geqslant 0$. Then

$$\begin{cases} w_t = w_{xx} - (\beta - h'(t))w_x + f(w), & t > 0,\ G(t) < x < 0, \\ w(t, G(t)) = 0,\ G'(t) = -\mu w_x(t, G(t)) + \mu w_x(t, 0), & t > 0, \\ w(t, 0) = 0,\ h'(t) = -\mu w_x(t, 0), & t > 0, \\ G(0) = -2h_0,\quad w(0, x) = u_0(x + h_0), & -2h_0 \leqslant x \leqslant 0. \end{cases} \tag{76}$$

It is easy to know that $G_\infty := \lim_{t\to\infty} G(t) = -\infty$. Since $w \in C^{1+\nu/2,2+\nu}(\Omega)$ with $\Omega := \{(t,x) \mid x \in [G(t),0],\ t \in [1,\infty)\}$, $h \in C^{1+\nu/2}([1,\infty))$ for any $\nu \in (0,1)$ and $h'(t) \to \beta - c_0$ by Lemma 4.13, there exists a sequence $\{t_n\}_{n=1}^\infty$ satisfying $t_n \to \infty$ as $n \to \infty$ such that

$$w(t+t_n,x) \to v(t,x) \text{ as } n \to \infty \text{ locally uniformly in } (t,x) \in \mathbb{R} \times (-\infty,0],$$

and v is a solution of

$$\begin{cases} v_t = v_{xx} - c_0 v_x + f(v), & t \in \mathbb{R},\ x < 0, \\ v(t,0) = 0,\ v_x(t,0) = -\frac{\beta - c_0}{\mu}, & t \in \mathbb{R}. \end{cases}$$

In case $\beta \in (c_0,\beta^*)$, we show that $v(t,x) \equiv V^*(x)$ for all $t \in \mathbb{R}$. If this is not true, then there exists $(t_0,x_0) \in \mathbb{R} \times (-\infty,0)$ such that $v(t_0,x_0) \neq V^*(x_0)$. Then for sufficiently small $\epsilon > 0$, when $t \in (0,\epsilon)$ we have $v(t_0+t,x_0) \neq V^*(x_0)$. Using zero number result Lemma 3.8 for $\eta(t,x) := v(t_0+t,x) - V^*(x)$ in $(t,x) \in [0,\epsilon] \times [x_0,0]$, we see that $\mathcal{Z}_{[x_0,0]}[\eta(t,\cdot)] < \infty$ for $t \in (0,\epsilon)$, and it decreases strictly once it has a degenerate point in $[x_0,0]$. This contradicts the fact that $x = 0$ is a degenerate zero of $\eta(t,\cdot)$ for all $t \in (0,\epsilon)$. Therefore, $v(t,x) \equiv V^*(x)$, and so $w(t+t_n,x) \to V^*(x)$ as $n \to \infty$ locally uniformly in $(t,x) \in \mathbb{R} \times (-\infty,0]$. By the uniqueness of $V^*(x)$ we actually prove $u(t,\cdot + h(t)) = w(t,\cdot) \to V^*(\cdot)$ as $t \to \infty$ uniformly in $[-M,0]$ for any $M > 0$.

In case $\beta = c_0$ we must have $v(t,x) \equiv 0$. Otherwise, $v(t,x) > 0$ for all $t \in \mathbb{R}$ and $x < 0$, and so the Hopf boundary lemma implies that $v_x(t,0) < 0$, contradicting $\beta = c_0$. Hence, $u(t,\cdot + h(t)) \to 0$ as $t \to \infty$ uniformly in $[-M,0]$ for any $M > 0$.

2. We prove the uniform convergence in $I(t)$ in case $c_0 < \beta < \beta^*$. For any small $\epsilon > 0$, there exists a large $M > 0$ such that

$$V^*(x) \leqslant V^*(-M) \leqslant \frac{\epsilon}{3} \text{ for } x \leqslant -M.$$

Taking $T > 0$ sufficiently large, by Step 1 we have

$$G(t) < -M, \quad \|u(t,\cdot + h(t)) - V^*(\cdot)\|_{L^\infty([-M,0])} < \frac{\epsilon}{3} \text{ for } t \geqslant T. \tag{77}$$

Hence, the function $u(t,\cdot + h(t))$ has a maximum point in $[-M,0]$. It is the unique maximum point by Lemma 4.14. Hence $u(t,\cdot + h(t))$ is increasing in $[G(t),-M]$, and so

$$0 \leqslant u(t,x+h(t)) \leqslant u(t,h(t)-M) \leqslant V^*(-M) + \frac{\epsilon}{3} \leqslant \frac{2\epsilon}{3} \quad \text{for } x \in [G(t),-M],\ t \geqslant T.$$

This implies that

$$\|u(t,\cdot + h(t)) - V^*(\cdot)\|_{L^\infty([G(t),-M])} \leqslant \epsilon \text{ for } t \geqslant T.$$

Combining with (77) we prove (74).

3. We now prove (75) in case $\beta = c_0$. By Lemma 4.14, $u(t,\cdot)$ has exactly one maximum point $\xi(t)$ when t is large, say, when $t \geqslant T$ for some $T > 0$. There are three cases:

Case 1. $u(t,\xi(t)) \to 0$ as $t \to \infty$;
Case 2. $u(t,\xi(t)) \to 1$ as $t \to \infty$;
Case 3. There exist $d \in (0,1)$ and a sequence $\{t_n\}_{n=1}^{\infty} \subset [T,\infty)$ with $t_n \to \infty$ such that $u(t_n,\xi(t_n)) = d$ for $n = 1,2,\cdots$.

The limit in (75) follows from Case 1 immediately. We now derive contradictions for Case 2 and Case 3.

Case 2: By Lemma 3.7, there exists $\delta_1 \in (0,c^*(\beta))$ such that the equation in (P) has a compactly supported traveling wave $W_{\delta_1}(x-\delta_1 t)$ with

$$W_{\delta_1}(0) = W_{\delta_1}(-L_{\delta_1}) = 0, \quad D_{\delta_1} := \max_{-L_{\delta_1}\leqslant z\leqslant 0} W_{\delta_1}(z) = \frac{1}{2} \quad \text{and} \quad \delta_1 = -\mu W'_{\delta_1}(0). \tag{78}$$

By Lemmas 4.1 and 4.4, $u(t,\cdot) \to 0$ as $t \to \infty$ uniformly in $[g(t), 2L_{\delta_1}]$, by the result in step 1 above, $u(t,\cdot) \to 0$ as $t \to \infty$ uniformly in $[h(t)-2L_{\delta_1}, h(t)]$. Hence we may assume that, for some $T_1 > T$,

$$2L_{\delta_1} < \xi(t) < h(t) - 2L_{\delta_1} \text{ and } u(t,\xi(t)) > D_{\delta_1} = \frac{1}{2} \text{ for all } t \geqslant T_1.$$

Now we consider the traveling wave $w_1(t,x) := W_{\delta_1}(x-\delta_1 t+\delta_1 T_1 - g_\infty)$. Clearly, when $t = T_1$ it has no contact point with $u(T_1,x)$. Since it moves rightward with speed $\delta_1 > 0$ and since $h'(t) \to 0$, the right endpoint $r_1(t) := \delta_1 t - \delta_1 T_1 + g_\infty$ of w_1 reaches $x = h(t)$ after some time. Before that, $r_1(t)$ first meets $g(t)$ at time $T_2 > T_1$, and then its left endpoint $l_1(t) := r_1(t) - L_{\delta_1}$ meets $g(t)$ at time $T_3 > T_2$. By Lemma 3.10, for $t \in [T_2, T_3)$ we have $\mathcal{Z}_{[g(t),r_1(t)]}[w_1(t,\cdot) - u(t,\cdot)] = 1$, and for $T_3 < t \ll T_3 + 1$, either

$$w_1(t,x) < u(t,x) \quad \text{for } x \in [l_1(t), r_1(t)], \tag{79}$$

or, $\mathcal{Z}_{[l_1(t),r_1(t)]}[w_1(t,\cdot) - u(t,\cdot)] = 2$. In the latter case, the two contact points between w_1 and u will not move across $x = \xi(t)$ where $u(t,\xi(t)) > \frac{1}{2} \geqslant w_1(t,\xi(t))$. Therefore, before $w_1(t,x)$ moves into the interval $[h(t) - L_{\delta_1}, h(t)]$, the two contact points disappear at some time $T_4 > T_3$, and so (79) holds for $t = T_4$. Once (79) holds at some time, it holds for all larger time since w_1 is a lower solution of (P). This leads to virtual spreading for u by Lemma 4.11, a contradiction.

Case 3: As above we select a compactly supported traveling wave $W_{\delta_2}(x-\delta_2 t)$ for some $\delta_2 \in (0,c^*(\beta))$ such that

$$W_{\delta_2}(0) = W_{\delta_2}(-L_{\delta_2}) = 0, \quad D_{\delta_2} := \max_{-L_{\delta_2}\leqslant z\leqslant 0} W_{\delta_2}(z) = W_{\delta_2}(-\tilde{z}) = d \quad \text{and}$$
$$\delta_2 = -\mu W'_{\delta_2}(0), \tag{80}$$

where $-\tilde{z} \in (-L_{\delta_2}, 0)$ is the maximum point of $W_{\delta_2}(z)$. By the locally uniform convergence in the above step 1 and in Lemma 4.1, there exists n_0 such that

$$2L_{\delta_2} < \xi(t_n) < h(t_n) - 2L_{\delta_2} \text{ for all } n \geqslant n_0. \tag{81}$$

Since $\xi(t_n) - \delta_2 t_n < h(t_n) - \delta_2 t_n \to -\infty$ as $n \to \infty$, there exists $n_1 > n_0$ such that

$$C := \xi(t_{n_1}) - \delta_2 t_{n_1} + \delta_2 t_{n_0} + \tilde{z} \leqslant g_\infty.$$

Now we consider the traveling wave $w_2(t,x) := W_{\delta_2}(x - \delta_2 t + \delta_2 t_{n_0} - C)$ for $t \geqslant t_{n_0}$. Since $w_2(t_{n_0}, x) = W_{\delta_2}(x - C)$, $w_2(t_{n_0}, \cdot)$ has no contact point with $u(t_{n_0}, x)$. Since w_2 moves rightward with speed $\delta_2 > 0$ and since $h'(t) \to 0$, the right endpoint $r_2(t) := \delta_2 t - \delta_2 t_{n_0} + C$ of w_2 reaches $x = h(t)$ after some time. Before that, $r_2(t)$ first meets $g(t)$ at some time $T_5 > t_{n_0}$, and then the left endpoint $l_2(t) := r_2(t) - L_{\delta_2}$ of w_2 meets $g(t)$ at some time $T_6 > T_5$. We remark that $T_6 < t_{n_1}$. In fact, by (81) we have

$$r_2(t_{n_1}) = \delta_2 t_{n_1} - \delta_2 t_{n_0} + C = \xi(t_{n_1}) + \tilde{z} > 2L_{\delta_2} > g(T_6) + L_{\delta_2} = r_2(T_6).$$

Now, for $t \in [T_5, T_6)$, using the zero number argument we have $\mathcal{Z}_{[g(t), r_2(t)]}[w_2(t,\cdot) - u(t,\cdot)] = 1$. For $T_6 < t \ll T_6 + 1$, we have either

$$w_2(t,x) < u(t,x) \quad \text{for } x \in [l_2(t), r_2(t)], \tag{82}$$

or, $\mathcal{Z}_{[l_2(t), r_2(t)]}[w_2(t,\cdot) - u(t,\cdot)] = 2$. (82) cannot be true, since it implies virtual spreading for u by Lemma 4.11. In case $w_2(t,\cdot) - u(t,\cdot)$ has two zeros for $T_6 < t \ll T_6 + 1$, by the zero number argument, the two zeros unite to be one degenerate zero $\xi(t_{n_1})$ at time t_{n_1} (note that $\xi(t_{n_1})$ is the maximum point of both $w_2(t_{n_1}, \cdot)$ and $u(t_{n_1}, \cdot)$). So after t_{n_1}, w_2 and u have no contact points. This implies that $w_2(t,x) < u(t,x)$ ($w_2 > u$ is impossible since the support of u is wider than that of w_2). This again leads to virtual spreading for u by Lemma 4.11, a contradiction.

This proves Theorem 4.15. □

Remark 4.16. By Lemma 4.13 we have $h(t) = (\beta - c_0)t + \varrho(t)$ for some $\varrho(t) = o(t)$. Hence the uniform convergence in (74) can be rewritten as (12).

4.4.4. Proofs of Theorems 2.2 and 2.3

In the last part of this subsection we prove Theorem 2.2 and Theorem 2.3. Remember we use $(u(t,x;\sigma\phi), g(t;\sigma\phi), h(t;\sigma\phi))$ to denote the solution of (P) with initial data $u_0 = \sigma\phi$ for some given $\phi \in \mathscr{X}(h_0)$. Define E_0 and σ_* as in (58), and when $c_0 \leqslant \beta < \beta^*$, denote

$$E_1 := \{\sigma > 0 \mid \text{virtual spreading happens for } (u,g,h)\}, \quad \sigma^* := \inf E_1.$$

By the comparison principle we have $[\sigma, \infty) \subset E_1$ if $\sigma \in E_1$. Thus $(\sigma^*, \infty) \subset E_1$.

Proof of Theorem 2.2. If $\sigma_* = \infty$, then there is nothing left to prove. We assume $\sigma_* \in (0, \infty)$ in the following.

We first prove $\sigma_* = \sigma^*$. Otherwise, $\sigma_* < \sigma^*$, and so there exist $\sigma_1, \sigma_2 \in (\sigma_*, \sigma^*)$ with $\sigma_1 < \sigma_2$. By the strong comparison principle we have

$$g(t; \sigma_1\phi) > g(t; \sigma_2\phi), \quad h(t; \sigma_1\phi) < h(t; \sigma_2\phi)$$

and

$$u(t, x; \sigma_1\phi) < u(t, x; \sigma_2\phi) \ \text{ for } x \in I^{\sigma_1}(t) := [g(t; \sigma_1\phi), h(t; \sigma_1\phi)], \ t > 0.$$

Since these inequalities are strict at $t = 1$, there exists $\epsilon > 0$ small such that

$$u(1, x; \sigma_1\phi) < u(1, x - \epsilon; \sigma_2\phi) \text{ for } x \in I^{\sigma_1}(1).$$

By the comparison principle again we have

$$u(t, x; \sigma_1\phi) < u(t, x - \epsilon; \sigma_2\phi) \text{ for } x \in I^{\sigma_1}(t), \ t \geqslant 1.$$

And so

$$\begin{aligned} &u(t, x + h(t; \sigma_1\phi); \sigma_1\phi) < u(t, x + h(t; \sigma_1\phi) - \epsilon; \sigma_2\phi) \\ &\quad \text{for } x \in [g(t; \sigma_1\phi) - h(t; \sigma_1\phi), 0], \ t \geqslant 1. \end{aligned} \tag{83}$$

By Theorem 4.15(i), both $u(t, x + h(t; \sigma_1\phi); \sigma_1\phi)$ and $u(t, x + h(t; \sigma_2\phi); \sigma_2\phi)$ converge to the tadpole-like function $V^*(x)$ uniformly. Taking limits as $t \to \infty$ in (83) we deduce a contradiction by $h(t; \sigma_1\phi) - \epsilon - h(t; \sigma_2\phi) \leqslant -\epsilon$. This proves $\sigma_* = \sigma^*$.

It is easily shown as in the proof of Theorem 4.9 that $E_0 \setminus \{0\}$ is open, and E_1 is open by Lemma 4.11, so neither vanishing nor virtual spreading happens for $(u(t, x; \sigma\phi), g(t; \sigma\phi), h(t; \sigma\phi))$ with $\sigma = \sigma^*$. Thus $u(t, x; \sigma^*\phi)$ is a transition solution and it converges to V^* as in Theorem 4.15 and Remark 4.16.

Other conclusions in Theorem 2.2 follow from the previous lemmas and theorems. □

Proof of Theorem 2.3. If $\sigma_* = \infty$, then there is nothing left to prove. If $\sigma_* \in (0, \infty)$ and $\sigma^* = \infty$, then vanishing happens for $u(t, x; \sigma\phi)$ with $\sigma < \sigma_*$, and virtual vanishing happens for $u(t, x; \sigma\phi)$ with $\sigma \geqslant \sigma_*$. Finally we consider the case $0 < \sigma_* \leqslant \sigma^* < \infty$. We show that E_1 is an open set. Indeed, if $\sigma_1 \in E_1$, then for any $\delta \in (0, c^*(\beta))$ there exist $T_1 > 0$, $x_1 \in \mathbb{R}$ such that

$$u(T_1, x; \sigma_1\phi) > W_\delta(x - x_1) \text{ for } x \in [x_1 - L_\delta, x_1];$$

since $u(T_1, \cdot\,; \sigma_1\phi)$ depends on σ_1 continuously, there exists $\epsilon > 0$ such that

$$u(T_1, x; \sigma\phi) > W_\delta(x - x_1) \text{ for } x \in [x_1 - L_\delta, x_1],$$

for any $\sigma \in [\sigma_1 - \epsilon, \sigma_1 + \epsilon]$. By Lemma 4.11, virtual spreading happens for $(u(t,x;\sigma\phi), g(t;\sigma\phi), h(t;\sigma\phi))$. Hence E_1 is an open set, and so $E_1 = (\sigma^*, \infty)$.

This proves the theorem. □

5. Uniform convergence when (virtual) spreading happens

In the main results Theorems 2.1, 2.2 and 2.3, we observe (virtual) spreading phenomena, which is the case where the solution converges to 1 locally uniformly in a fixed or moving coordinate frame. In this section we consider the asymptotic profiles for such solutions in the whole domain.

Throughout this section we assume $0 < \beta < \beta^*$.

5.1. Locally uniform convergence of the front

We first describe the asymptotic profile near the front $x = h(t)$. In a similar way as [16,26,29] one can show that:

Proposition 5.1. *Assume $0 < \beta < \beta^*$. If (virtual) spreading happens for a solution of (P), then there exists $H_\infty \in \mathbb{R}$ such that*

$$\lim_{t\to\infty}[h(t) - c^*t] = H_\infty, \ \lim_{t\to\infty} h'(t) = c^*, \tag{84}$$

$$\lim_{t\to\infty} u(t, \cdot + h(t)) = U^*(\cdot) \text{ locally uniformly in } (-\infty, 0]. \tag{85}$$

For small advection: $0 < \beta < c_0$, one can give a uniform convergence for the solution (u, g, h) of (P) as in [16,26,29].

Proposition 5.2. *Assume $0 < \beta < c_0$. If spreading happens for a solution (u, g, h) of (P), then there exist $G_\infty, H_\infty \in \mathbb{R}$ such that (84) holds and*

$$\lim_{t\to\infty}[g(t) - c_l^*t] = G_\infty, \ \lim_{t\to\infty} g'(t) = c_l^*,$$

$$\lim_{t\to\infty} \|u(t,\cdot) - U^*(\cdot - c^*t - H_\infty)\cdot U_l^*(\cdot - c_l^*t - G_\infty)\|_{L^\infty([g(t),h(t)])} = 0,$$

if we extend U^* and U_l^* to be zero outside their supports.

5.2. Locally uniform convergence of the back

In this subsection we show that, when $c_0 \leqslant \beta < \beta^*$, the back of a virtual spreading solution u converges to a traveling wave Q locally uniformly. We will use the following definition:

Definition 5.3. (See [17].) Let u_1, u_2 be two entire solutions of $u_t = u_{xx} - \beta u_x + f(u)$ satisfying $u_{1x}(t,x) > 0$ and $u_{2x}(t,x) > 0$ for all $x \in \mathbb{R}, t \in \mathbb{R}$. We say that u_1 is **steeper than** u_2 if for any t_1, t_2 and x_1 in $\mathbb{R}$ such that $u_1(t_1,x_1) = u_2(t_2,x_1)$, we have either

$$u_1(\cdot + t_1, \cdot) \equiv u_2(\cdot + t_2, \cdot) \text{ or } (u_1)_x(t_1,x_1) > (u_2)_x(t_2,x_1).$$

As above, u_1 and u_2 are called entire solutions since they are defined for all $t \in \mathbb{R}$. The above property implies that the graph of the solution u_1 (at any chosen time moment t_1) and that of the solution u_2 (at any chosen time moment t_2) can intersect at most once unless they are identical, and that if they intersect at a single point, then $u_1 - u_2$ is negative on the left-hand side of the intersection point, while positive on the right-hand side.

Theorem 5.4. *Assume $\beta \in [c_0, \beta^*)$. If virtual spreading happens for a solution (u,g,h) of (P), then there exists a continuous function $\theta(t)$ with $\theta(t) = o(t)$ and $\theta(t) \to \infty$ $(t \to \infty)$ such that for any $M > 0$,*

$$\lim_{t\to\infty} \|u(t,\cdot) - Q(\cdot - (\beta - c_0)t - \theta(t))\|_{L^\infty([g(t),(\beta-c_0)t+\theta(t)+M])} = 0. \tag{86}$$

Proof. The proof is long and is divided into several steps. We will use C and T to denote positive constants which may be different case by case.

Step 1. A rough estimate for the speed of the back. For any $\delta_1 \in (0, c^*(\beta) - \beta + c_0)$, we consider the compactly supported traveling wave $W_{\delta_1}(x - c_1 t)$ with $c_1 = \beta - c_0 + \delta_1$, where $W_{\delta_1}(z) := W(z; c_1, -c_0 + \delta_1)$ is the solution of (23) and (32) with $c = c_1$, whose support is $[-L_{\delta_1}, 0] = [-L(c_1, -c_0 + \delta_1), 0]$. As in Lemma 3.7 we denote $D_{\delta_1} := \max_{-L_{\delta_1} \leqslant z \leqslant 0} W_{\delta_1}(z)$. Write

$$m := \frac{1}{2} \inf\{D_\delta \mid 0 < \delta < c^*(\beta) - \beta + c_0\}. \tag{87}$$

Then $m \in (0, \frac{1}{2})$ by the phase plane analysis.

By our assumption, virtual spreading happens: $u(t, \cdot + ct) \to 1$ locally uniformly in $\mathbb{R}$ for some $c > 0$. Hence for any given $\delta_1 \in (0, c^* - \beta + c_0)$ there exist a large T_0 and $r \in \mathbb{R}$ such that

$$u(T_0, x) \geqslant W_{\delta_1}(x - r) \quad \text{for } x \in [r - L_{\delta_1}, r].$$

By comparison we have

$$u(t + T_0, x) > W_{\delta_1}(x - c_1 t - r) \quad \text{for } x \in [r + c_1 t - L_{\delta_1}, r + c_1 t],\ t > 0.$$

Therefore $\chi(t) := \min\{x \in [g(t), h(t)] \mid u(t,x) = m\}$ satisfies

$$\chi(t + T_0) < c_1 t + r \quad \text{for } t > 0. \tag{88}$$

Combining with (50) we have

$$(\beta - c_0)t + \frac{3}{c_0}\ln t - C \leqslant \chi(t) \leqslant (\beta - c_0 + \delta_1)t + C, \quad t \gg 1. \tag{89}$$

Step 2. Truncation of the solution. Instead of u we will consider its truncation on $[g(t), \xi(t)]$ for some $\xi(t) \in (g(t), h(t))$.

Let $\epsilon \in (0, \frac{1}{2}(1-m))$ be any given small constant. We define $\xi(t) < h(t)$ as a position closest to $h(t)$ where u takes value $1-\epsilon$. More precisely, by the definition of the rightward traveling semi-wave $U^*(x - c^*(\beta)t)$, there exists $M_1 = M_1(\epsilon) > 0$ sufficiently large such that $U^*(-M_1) > 1 - \frac{\epsilon}{2}$. By (85), $u(t, h(t) - M_1) > 1 - \epsilon$ for sufficiently large t, and so there exists $\xi(t) \in [h(t) - M_1, h(t)]$ such that

$$u(t, \xi(t)) = 1 - \epsilon, \quad \text{for large } t. \tag{90}$$

By (89) and (84) we have

$$\xi(t) - \chi(t) \geqslant h(t) - M_1 - \chi(t) \geqslant (c^* - \beta + c_0 - \delta_1)t + O(1) \to \infty, \quad \text{as } t \to \infty. \tag{91}$$

Since the convergence in (85) holds in fact in $C^2_{\text{loc}}((-\infty, 0])$ topology by parabolic estimate, it follows from $U^*_x(x) < 0$ that

$$u_x(t, x) < 0 \text{ for } x \in [h(t) - 2M_1, h(t)] \text{ and } t \gg 1. \tag{92}$$

So the leftmost local maximum point $\xi_1(t)$ of $u(t, \cdot)$ satisfies $\xi_1(t) < h(t) - 2M_1 < \xi(t)$ for $t \gg 1$.

We now show that

$$u(t, x) \geqslant 1 - \epsilon \text{ for } x \in [\xi_1(t), \xi(t)], \ t \gg 1. \tag{93}$$

In case $u(t, \cdot)$ has exactly one local maximum point $\xi_1(t)$ for large t, (93) holds since $u(t, \cdot)$ is decreasing in $[\xi_1(t), \xi(t)]$ and $u(t, \xi(t)) = 1-\epsilon$. We now consider the case that $u(t, \cdot)$ has exactly $N\,(\geqslant 2)$ local maximum points $\{\xi_i(t)\}_{i=1}^N$ with $g(t) < \xi_1(t) < \cdots < \xi_N(t) < h(t)$ for large t. We remark that this case is possible only if $\beta < c^*(\beta)$. In fact, when $\beta \geqslant c^*(\beta)$, by (37) and (84) we have

$$0 < h(t) - \xi_1(t) < (c^*(\beta) - \beta)t + C \leqslant C, \quad t \gg 1.$$

This contradicts the locally uniform convergence (85) and the fact that U^* is a strictly decreasing function. So, in the following, we assume that

$$\beta < c^*(\beta) \quad \text{and} \quad \xi_1(t) \geqslant \beta t - C \text{ for some } C > 0. \tag{94}$$

Choose a small $\delta \in (0, c_0)$ and consider the solution $q(z) := W(z; b, -\delta)$ of (20) with $\gamma = -\delta$, where $b \in (0, P(-\delta))$. This solution corresponds to a point $G \in S_1$ as in Fig. 2(a), and its trajectory is a curve like Γ_2 in Fig. 1(a). When $b \to P(-\delta)$, the trajectory $\Gamma_2 \to \Gamma_1$ in Fig. 1(a). As in Section 3.2, for the above given $\epsilon > 0$ and $\delta \in (0, c_0)$, there exists $b = b(\epsilon, \delta) \in (0, P(-\delta))$ such that $\mathscr{W}(z) := W(z; b(\epsilon, \delta), -\delta)$ and $\mathcal{L} = L(b(\epsilon, \delta), -\delta)$ satisfy

$$\mathscr{W}(0) = \mathscr{W}(-\mathcal{L}) = 0, \quad \mathscr{W}(z) > 0 \text{ for } z \in (-\mathcal{L}, 0),$$
$$\mathscr{W}(\hat{z}) = \max_{-\mathcal{L} \leqslant z \leqslant 0} \mathscr{W}(z) = 1 - \epsilon \quad \text{for some } \hat{z} \in (-\mathcal{L}, 0).$$

We prove (93) by contradiction. By (92) we only need to prove (93) for $x \in [\xi_1(t), h(t) - 2M_1]$. Assume that there exist a time sequence $\{t_n\}_{n=1}^{\infty}$ and a sequence $\{y_n\}_{n=1}^{\infty}$ with $t_n \to \infty$ and $y_n \in [\xi_1(t_n), h(t_n) - 2M_1]$ such that

$$u(t_n, y_n) < 1 - \epsilon \text{ for all } n. \tag{95}$$

For each n, we define a continuous function of τ by

$$\psi_n(\tau) := h(\tau) + (\beta - \delta)(t_n - \tau) - y_n - M_1 + \hat{z}.$$

It is easily seen that

$$\psi_n(t_n) \geqslant h(t_n) - (h(t_n) - 2M_1) - M_1 + \hat{z} > 0.$$

For $\rho \in (0, 1)$, by $y_n \geqslant \xi_1(t_n) \geqslant \beta t_n - C$ we have

$$\begin{aligned}\psi_n(\rho t_n) &= h(\rho t_n) + (\beta - \delta)(1 - \rho)t_n - y_n + O(1) \\ &\leqslant [c^* \rho - (\beta - \delta)\rho - \delta]t_n + O(1) \to -\infty \text{ as } n \to \infty,\end{aligned}$$

provided $\rho > 0$ is sufficiently small. Hence for such a ρ and for any large n, there exists $\tau_n \in (\rho t_n, t_n)$ such that $\psi_n(\tau_n) = 0$.

For any large n, by (85) we have

$$\begin{aligned}u(\tau_n, x) &\geqslant U^*(x - h(\tau_n)) - \frac{\epsilon}{2} \\ &\geqslant \mathscr{W}(x - h(\tau_n) + M_1), \quad x \in [h(\tau_n) - M_1 - \mathcal{L}, h(\tau_n) - M_1].\end{aligned}$$

Set $r_n(t) := (\beta - \delta)t + h(\tau_n) - M_1$. Since $\mathscr{W}(x - r_n(t))$ is a compactly supported traveling wave of $(P)_1$, and its right endpoint

$$r_n(t) = (\beta - \delta)t + c^* \tau_n + H_\infty - M_1 + o(1) < h(t + \tau_n) = c^* t + c^* \tau_n + H_\infty + o(1)$$

by (84) and (94), provided n is sufficiently large. Hence $\mathscr{W}(x - r_n(t))$ is a lower solution of (P) and by the comparison principle we have

$$u(t + \tau_n, x) \geqslant \mathscr{W}(x - r_n(t)) \ \text{ for } x \in [r_n(t) - \mathcal{L}, r_n(t)], \ t > 0.$$

In particular, at $t = t_n - \tau_n > 0$ and $x = y_n$, by $\psi_n(\tau_n) = 0$ we have

$$1 - \epsilon > u(t_n, y_n) \geqslant \mathscr{W}(\hat{z}) = 1 - \epsilon,$$

a contradiction. This proves (93).

In what follows, we write $\hat{u}(t, x) := u(t, x)\big|_{x \in [g(t), \xi(t)]}$ as a truncation of u.

Step 3. Truncation of tadpole-like traveling waves. For any $b \in (0, P(-c_0))$ recall that $V(z; b, -c_0)$ is a tadpole-like solution of (20) (cf. point H in Fig. 2(a)). We choose $b = b(\epsilon)$ near $P(-c_0)$ such that

$$\max_{z \leqslant 0} V(z; b(\epsilon), -c_0) = V(\bar{z}; b(\epsilon), -c_0) = 1 - 2\epsilon > m,$$

for some $\bar{z} < 0$. In a similar way as above, we write

$$\widehat{V}(x - (\beta - c_0)t) := V(x - (\beta - c_0)t; b(\epsilon), -c_0)\big|_{x \in (-\infty, \bar{z} + (\beta - c_0)t]}$$

as a truncation V.

Step 4. Comparison between $\hat{u}$ and $\widehat{V}$. To study the asymptotic profile of $\hat{u}$, we compare $\hat{u}$ with a family of the shifts of $\widehat{V}$. Without loss of generality, we may assume $u(t, x)$ satisfies all the properties in Steps 1–3 from time $t = 0$. Since $\hat{u}_x(0, g(0)) > 0$, one can choose $X > 0$ large such that $\hat{u}(0, x)$ and $\widehat{V}(x - \hat{x})$ (for any $\hat{x} \geqslant X$) intersect at exactly one point $\hat{y}$, and

$$\hat{u}(0, x) < \widehat{V}(x - \hat{x}) \text{ for } x \in [g(0), \hat{y}), \ \hat{u}(0, x) > \widehat{V}(x - \hat{x}) \text{ for } x \in (\hat{y}, \min\{\xi(0), \bar{z} + \hat{x}\}).$$

Since the back of $u(t, \cdot)$ moves rightward faster than $(\beta - c_0)t + \frac{3}{c_0} \ln t - C$ by (50), it will exceed $\widehat{V}(x - (\beta - c_0)t - \hat{x})$ at some time $\hat{T} > 0$, that is, their intersection point $(y(t), \hat{u}(t, y(t)))$ starting from $(\hat{y}, \hat{u}(0, \hat{y}))$ exists only in time interval $[0, \hat{T}]$.

For each $\hat{x} \geqslant X$, both $\hat{u}(t, x)$ and $\widehat{V}(x - (\beta - c_0)t - \hat{x})$ are solutions of $(P)_1$. We now compare them and show that for $t \in [0, \hat{T})$,

$$\begin{cases} \text{there exists } y(t) \in (g(t), \xi(t)) \cap (-\infty, \eta(t)] \text{ such that} \\ \quad \hat{u}(t, x) < \widehat{V}(x - (\beta - c_0)t - \hat{x}) \text{ for } x \in [g(t), y(t)), \\ \quad \hat{u}(t, x) > \widehat{V}(x - (\beta - c_0)t - \hat{x}) \text{ for } x \in \big(y(t), \min\{\xi(t), \eta(t)\}\big], \end{cases} \tag{96}$$

where $\eta(t) := (\beta - c_0)t + \hat{x} + \bar{z}$. By the comparison principle, the conclusions in (96) can be proved if we can exclude the following two possibilities:

(A) the right endpoint $(\xi(t), 1-\epsilon)$ of $\hat{u}(t,\cdot)$ touches $\widehat{V}$ at some time $t \in (0, \hat{T})$;
(B) the right endpoint $(\eta(t), 1-2\epsilon)$ of $\widehat{V}(x-(\beta-c_0)t-\hat{x})$ touches $\hat{u}$ at some time $t \in (0, \hat{T})$.

(A) of course is impossible because $\hat{u}(t,\cdot)$ takes value $1-\epsilon$ at $x=\xi(t)$, bigger than $\max \widehat{V}$. (B) is impossible when $\eta(t) \in [\xi_1(t), \xi(t)]$ since in this case $\hat{u}(t,\eta(t)) \geqslant 1-\epsilon > \max \widehat{V}$ by (93). When $\eta(t) < \xi_1(t)$, $\widehat{V}_x(x-(\beta-c_0)t-\hat{x})\big|_{x=\eta(t)} = \widehat{V}_x(\bar{z}) = 0$ and $\hat{u}_x(t,\eta(t)) > 0$. Hence $(\eta(t), 1-2\epsilon)$ cannot be a new emerging intersection point between $\hat{u}$ and $\widehat{V}$. This excludes the possibility of (B).

Step 5. Slope of the back of $u(t,\cdot)$. By (96), there exists exactly one intersection point $(y(t), \hat{u}(t,y(t)))$ between $\hat{u}(t,x)$ and $\widehat{V}(x-(\beta-c_0)t-\hat{x})$. Moreover, the zero number argument implies that this point is non-degenerate for $t \in (0, \hat{T})$. Hence

$$\hat{u}(t,y(t)) = \widehat{V}(y(t)-(\beta-c_0)t-\hat{x}) \quad \text{and}$$
$$\hat{u}_x(t,y(t)) > \widehat{V}_x(y(t)-(\beta-c_0)t-\hat{x}), \quad t \in (0,\hat{T}).$$

Since $\widehat{V}(z)$ is strictly increasing in $(-\infty, \bar{z}]$ (cf. Section 3.2) and since $\widehat{V}(\bar{z}) = 1-2\epsilon > m$, the equation $\widehat{V}(z) = m$ has a unique root y^0 in $(-\infty, \bar{z})$. For any given large t, we take $\hat{x} = \chi(t)-(\beta-c_0)t-y^0$, then the function $\hat{u}(t,x)$ and $\widehat{V}(x-(\beta-c_0)t-\hat{x})$ intersect exactly at $x=\chi(t)$:

$$\hat{u}(t,\chi(t)) = m = \widehat{V}(y^0) = \widehat{V}(\chi(t)-(\beta-c_0)t-\hat{x}).$$

By the Hopf lemma we have

$$\hat{u}_x(t,\chi(t)) > \widehat{V}_x(y^0). \tag{97}$$

Step 6. Convergence of the back of u and the slope of the limit function. For any increasing sequence $\{t_n\}_{n=0}^{\infty}$ with $t_n \to \infty$ $(n \to \infty)$, we set $x_n := \chi(t_n)$ and define

$$\hat{u}_n(t,x) := \hat{u}(t+t_n, x+x_n) \text{ for } g(t+t_n)-x_n \leqslant x \leqslant \xi(t+t_n)-x_n,\ t > -t_n.$$

Clearly, $\hat{u}_n(0,0) = \hat{u}(t_n,x_n) = m$ for $n \in \mathbb{N}$. For any given $t \in \mathbb{R}$, $g(t+t_n)-x_n \to -\infty$ as $n \to \infty$ and by (84) and (91) we have

$$\begin{aligned}\xi(t+t_n)-x_n &= \xi(t+t_n)-\xi(t_n)+[\xi(t_n)-\chi(t_n)]\\ &\geqslant h(t+t_n)-M_1-h(t_n)+[\xi(t_n)-\chi(t_n)]\\ &= c^*t+O(1)+[\xi(t_n)-\chi(t_n)] \to \infty \text{ as } n \to \infty.\end{aligned}$$

Since $\hat{u}_n(t,x)$ is bounded in L^∞ norm, by parabolic estimate, it is also bounded in $C^{1+\nu/2,2+\nu}([-M,M]\times[-M,M])$ norm for any $M>0$ and any $\nu \in (0,1)$. By Cantor's diagonal argument, there exists a subsequence $\{n_j\}$ of $\{n\}$ such that

$$\lim_{j\to\infty} \hat{u}_{n_j}(t,x) = w(t,x) \text{ in } C^{1,2}_{\mathrm{loc}}(\mathbb{R}^2) \text{ topology},$$

where $w \in C^{1,2}(\mathbb{R}^2)$ is an entire solution of $(P)_1$ with $w(0,0) = m$. By (97) we have

$$w_x(0,0) = \lim_{j\to\infty} (\hat{u}_{n_j})_x(0,0) = \lim_{j\to\infty} \hat{u}_x(t_{n_j}, x_{n_j}) = \lim_{j\to\infty} \hat{u}_x(t_{n_j}, \chi(t_{n_j})) \geqslant \widehat{V}_x(y^0). \quad (98)$$

For the solution $Q(z)$ of (23)–(24) with $c = \beta - c_0$, there exists a unique $y^* \in \mathbb{R}$ such that $Q(y^*) = m$. By the phase plane analysis (Lemma 3.5(i)), $V(\cdot + y^0; b(\epsilon), -c_0) = \widehat{V}(\cdot + y^0) \to Q(\cdot + y^*)$ in $C^2_{\mathrm{loc}}(\mathbb{R})$ topology, as $\epsilon \to 0$, or equivalently, as $b(\epsilon) \to P(-c_0)$. Taking limit as $\epsilon \to 0$ in (98) we have

$$w_x(0,0) \geqslant Q'(y^*). \quad (99)$$

On the other hand, both $u_1(t,x) := Q(x - (\beta - c_0)t + y^*)$ and $u_2(t,x) := w(t,x)$ are entire solutions of $(P)_1$. By [17, Lemma 2.8], u_1 is steeper than u_2 in the sense of Definition 5.3. In particular, taking $t_1 = t_2 = x_1 = 0$ in Definition 5.3 we have $u_1(0,0) = Q(y^*) = m = w(0,0) = u_2(0,0)$. Hence

$$w(t,x) \equiv Q(x - (\beta - c_0)t + y^*) \quad \text{for all } t, x \in \mathbb{R}$$

by Definition 5.3 and the inequality (99).

Therefore, $\lim_{j\to\infty} \hat{u}_{n_j}(t,x) = Q(x - (\beta - c_0)t + y^*)$ in $C^{1,2}_{\mathrm{loc}}(\mathbb{R}^2)$ topology. By the uniqueness of the limit function Q we have

$$\lim_{n\to\infty} u_n(t,x) = \lim_{n\to\infty} u(t + t_n, x + \chi(t_n)) = Q(x - (\beta - c_0)t + y^*) \text{ in } C^{1,2}_{\mathrm{loc}}(\mathbb{R}^2) \text{ topology}.$$

Since $\{t_n\}$ is an arbitrarily chosen sequence we obtain

$$\lim_{\tau\to\infty} u(t + \tau, x + \chi(\tau)) = Q(x - (\beta - c_0)t + y^*) \text{ in } C^{1,2}_{\mathrm{loc}}(\mathbb{R}^2) \text{ topology}.$$

Taking $t = 0$ we have

$$\lim_{\tau\to\infty} u(\tau, x + \chi(\tau)) = Q(x + y^*) \text{ in } C^2_{\mathrm{loc}}(\mathbb{R}) \text{ topology}. \quad (100)$$

Define

$$\theta(\tau) := \chi(\tau) - (\beta - c_0)\tau - y^*,$$

which is a continuous function of τ. Then by (89) we have

$$\frac{3}{c_0} \ln \tau - C \leqslant \theta(\tau) \leqslant \delta_1 \tau + C, \quad \tau \gg 1.$$

Since this is true for any small $\delta_1 > 0$ (see Step 1) we have $\theta(\tau) = o(\tau)$ $(\tau \to \infty)$. Thus by (100) we have

$$\lim_{\tau\to\infty} [u(\tau, x) - Q(x - (\beta - c_0)\tau - \theta(\tau))] = 0 \tag{101}$$

uniformly in $[(\beta - c_0)\tau + \theta(\tau) - M, (\beta - c_0)\tau + \theta(\tau) + M]$ for any $M > 0$.

Step 7. Completion of the proof. For any $\varepsilon_0 > 0$, there exists $M > 0$ such that $Q(z) \leqslant Q(-M) \leqslant \varepsilon_0$ for $z < -M$. For this M, we choose $T > 0$ large such that when $t > T$ we have

$$u(t, (\beta - c_0)t + \theta(t) - M) \leqslant 2Q(-M) \leqslant 2\varepsilon_0$$

by (101). $u(t, \cdot)$ is increasing in $[g(t), (\beta - c_0)t + \theta(t) - M]$ by Lemma 3.11, hence, when $t > T$ and $x \in [g(t), (\beta - c_0)t + \theta(t) - M]$ we have

$$|u(t, x) - Q(x - (\beta - c_0)t - \theta(t))| \leqslant u(t, (\beta - c_0)t + \theta(t) - M) + Q(-M) \leqslant 3\varepsilon_0.$$

Combining with (101) we obtain the conclusion (86). □

5.3. Uniform convergence

In this subsection, we complete the proof of Theorem 2.5.

Proof of Theorem 2.5. (15), (16) and (17) are proved in Propositions 5.1 and 5.2. We only need to prove (18) for $c_0 \leqslant \beta < \beta^*$. For any given small $\varepsilon > 0$, we will prove

$$\begin{aligned} &|u(t, x) - U^*(x - c^*t - H_\infty) \cdot Q(x - (\beta - c_0)t - \theta(t))| \\ &\quad < C\varepsilon \quad \text{for } x \in I(t) \text{ and large } t, \end{aligned} \tag{102}$$

where $C > 0$ is a constant independent of t and x.

By Lemma 3.7, for any $\delta \in (0, c^*(\beta) - \beta + c_0)$ the problem (P) has a compactly supported traveling wave $W_\delta(x - (\beta - c_0 + \delta)t)$, where $W_\delta(z)$ (with support $[-L_\delta, 0]$) is the unique solution of the problems (23) and (32), whose maximum and maximum point are denoted by D_δ and $-z_\delta$, respectively. Moreover, for the above given $\varepsilon > 0$, Lemma 3.7 also indicates that, there exists $\delta_\varepsilon \in (0, c^*(\beta) - \beta + c_0)$ such that

$$D_\delta = W_\delta(-z_\delta) \in (1 - \varepsilon, 1) \text{ when } \delta \in (\delta_\varepsilon, c^*(\beta) - \beta + c_0).$$

We select $\delta_1, \delta_0, \delta_2 \in (\delta_\varepsilon, c^*(\beta) - \beta + c_0)$ with $\delta_1 > \delta_0 > \delta_2$ and fix them. For $i = 1$ and 2, denote $\kappa_i = (1 - D_{\delta_i})/(3\varepsilon)$, then $\kappa_i \in (0, \frac{1}{3})$.

By the definitions of $U^*(z)$ and $Q(z)$, there exists $M(\varepsilon) > 0$ such that when $M > M(\varepsilon)$,

$$1-\varepsilon \leqslant U^*(z) \leqslant 1 \text{ for } z \leqslant -M, \qquad 1-\varepsilon \leqslant Q(z) \leqslant 1 \text{ for } z \geqslant M \tag{103}$$

and there exists $M(\delta_1, \delta_2) > M(\varepsilon)$ such that when $M > M(\delta_1, \delta_2)$,

$$\begin{aligned} Q(z) &> D_{\delta_1} + \kappa_1 \varepsilon \text{ for } z \in [M - L_{\delta_1}, M], \\ U^*(z) &> D_{\delta_2} + \kappa_2 \varepsilon \text{ for } z \in [-2M, -2M + L_{\delta_2}]. \end{aligned} \tag{104}$$

In what follows we fix an $M > M(\delta_1, \delta_2) > M(\varepsilon)$.

Since the solution $\eta(t)$ of the problem

$$\eta_t = f(\eta), \quad \eta(0) = 1 + \|u_0\|_{L^\infty}$$

is an upper solution of (P) and since $\eta(t) \to 1$ as $t \to \infty$, there exists a time $T_1 = T_1(\varepsilon) > 0$ such that

$$u(t,x) < 1 + \varepsilon \quad \text{for } x \in I(t),\ t > T_1. \tag{105}$$

By Theorem 5.4, there exists $T_2 > T_1$ such that when $t > T_2$ we have

$$\begin{aligned} &|u(t,x) - Q(x - (\beta - c_0)t - \theta(t))| < \kappa_1 \varepsilon \\ &\quad \text{for } x \in I_l(t) := [g(t), (\beta - c_0)t + \theta(t) + M], \end{aligned} \tag{106}$$

where $\theta(t)$ is a continuous positive function with $\theta(t) = o(t)$ and $\theta(t) \to \infty$ $(t \to \infty)$. By Proposition 5.1, there exists $T_3 > T_2$ such that when $t > T_3$ we have

$$|u(t,x) - U^*(x - c^* t - H_\infty)| < \kappa_2 \varepsilon \text{ for } x \in I_r(t) := [h(t) - 2M, h(t)] \tag{107}$$

(we extend $U^*(z)$ to be zero for $z > 0$ if necessary). We now prove

$$u(t,x) \geqslant 1 - \varepsilon \text{ for } x \in I_c(t) := [(\beta - c_0)t + \theta(t) + M, h(t) - 2M] \text{ and large } t. \tag{108}$$

Once this is proved, combining it with the above results we obtain (102) with $C = 3$.

In the following we prove (108) by contradiction. Assume that there exist a time sequence $\{t_n\}_{n=1}^\infty$ with $t_n \to \infty$ and a sequence $\{y_n\}$ with $y_n \in I_c(t_n)$ for each n such that

$$u(t_n, y_n) < 1 - \varepsilon \quad \text{for all } n. \tag{109}$$

We divide the interval $I_c(t)$ into $I_c^1(t)$ and $I_c^2(t)$, where

$$I_c^1(t) := [(\beta - c_0)t + \theta(t) + M, (\beta - c_0 + \delta_0)t], \quad I_c^2(t) := [(\beta - c_0 + \delta_0)t, h(t) - 2M].$$

Our idea to derive contradictions is the following. We put a compactly supported traveling wave $W_{\delta_1}(x - (\beta - c_0 + \delta_1)t + C_1)$ (resp. $W_{\delta_2}(x - (\beta - c_0 + \delta_2)t + C_2)$) under

$u(t+\tau,x)$ at time $t=0$ in the interval $I_l(\tau)$ (resp. $I_r(\tau)$), and then as t increases to $t_n-\tau$, its maximum point exactly reaches $y_n \in I_c^1(t_n)$ (resp. $y_n \in I_c^2(t_n)$), this leads to a contradiction.

First we consider the case that $\{y_n\}$ has a subsequence (denote it again by $\{y_n\}$) such that $y_n \in I_c^1(t_n)$ for each n. Define a continuous function of τ:

$$\psi_n^{(1)}(\tau) := \delta_1\tau - \theta(\tau) + y_n - (\beta - c_0 + \delta_1)t_n - M + z_{\delta_1} \text{ for } \tau \leqslant t_n.$$

Since $y_n \in I_c^1(t_n)$, it is easily seen that

$$\psi_n^{(1)}(t_n) = y_n - (\beta - c_0)t_n - \theta(t_n) - M + z_{\delta_1} \geqslant z_{\delta_1} > 0,$$

and for $\rho_1 = \frac{1}{2}(1 - \frac{\delta_0}{\delta_1}) \in (0,1)$ we have

$$\psi_n^{(1)}(\rho_1 t_n) \leqslant (\rho_1\delta_1 + \delta_0 - \delta_1)t_n + O(1) \to -\infty \text{ as } n \to \infty.$$

Hence, when n is sufficiently large, there exists $\tau_n \in (\rho_1 t_n, t_n)$ such that $\psi_n^{(1)}(\tau_n) = 0$. For such a large n, by (106) and (104) we have

$$u(\tau_n, x) \geqslant Q(x - (\beta - c_0)\tau_n - \theta(\tau_n)) - \kappa_1\varepsilon \geqslant D_{\delta_1} \geqslant W_{\delta_1}(x - X) \quad \text{for } x \in [X - L_{\delta_1}, X],$$

where $X := (\beta - c_0)\tau_n + \theta(\tau_n) + M$. Using comparison principle we have

$$u(t+\tau_n, x) \geqslant W_{\delta_1}(x - (\beta - c_0 + \delta_1)t - X) \text{ for } x \in J_1(t),\ t > 0,$$

where $J_1(t) := [(\beta - c_0 + \delta_1)t + X - L_{\delta_1}, (\beta - c_0 + \delta_1)t + X]$. By $\psi_n^{(1)}(\tau_n) = 0$ we have

$$y_n = (\beta - c_0 + \delta_1)(t_n - \tau_n) + X - z_{\delta_1} \in J_1(t_n - \tau_n).$$

Hence by taking $t = t_n - \tau_n$ and $x = y_n$ we have

$$1 - \varepsilon > u(t_n, y_n) \geqslant W_{\delta_1}(y_n - (\beta - c_0 + \delta_1)(t_n - \tau_n) - X) = W_{\delta_1}(-z_{\delta_1}) = D_{\delta_1} > 1 - \varepsilon,$$

a contradiction.

Next we consider the case that $\{y_n\}$ has a subsequence (denote it again by $\{y_n\}$) such that $y_n \in I_c^2(t_n)$ for each n. The proof is similar as above. Define a continuous function

$$\psi_n^{(2)}(\tau) := h(\tau) - (\beta - c_0 + \delta_2)\tau - 2M + L_{\delta_2} - y_n + (\beta - c_0 + \delta_2)t_n - z_{\delta_2} \text{ for } \tau \leqslant t_n.$$

Since $y_n \in I_c^2(t_n)$, it is easily seen that

$$\psi_n^{(2)}(t_n) = h(t_n) - 2M + L_{\delta_2} - y_n - z_{\delta_2} \geqslant L_{\delta_2} - z_{\delta_2} > 0,$$

and for $\rho_2 \in (0,1)$ we have

$$\begin{aligned}\psi_n^{(2)}(\rho_2 t_n) &\leqslant h(\rho_2 t_n) - (\beta - c_0 + \delta_2)\rho_2 t_n - y_n + (\beta - c_0 + \delta_2)t_n + O(1)\\ &= c^*(\beta)\rho_2 t_n + (\beta - c_0 + \delta_2)(1-\rho_2)t_n - y_n + O(1)\\ &\leqslant [-\rho_2(\beta - c_0 + \delta_2) + c^*(\beta)\rho_2 - \delta_0 + \delta_2]t_n + O(1).\end{aligned}$$

Since $\delta_0 > \delta_2$, the coefficient of t_n in the last line is negative when $\rho_2 > 0$ is sufficiently small. Hence for such a ρ_2, $\psi_n^{(2)}(\rho_2 t_n) \to -\infty$ as $n \to \infty$. Consequently, for any large n, there exists $\tau_n' \in (\rho_2 t_n, t_n)$ such that $\psi_n^{(2)}(\tau_n') = 0$. By (107) and (104) we have

$$u(\tau_n', x) \geqslant U^*(x - c^*\tau_n' - H_\infty) - \kappa_2\varepsilon > D_{\delta_2} \geqslant W_{\delta_2}(x - X') \quad \text{for } x \in [X' - L_{\delta_2}, X'],$$

where $X' := h(\tau_n') - 2M + L_{\delta_2}$. By the comparison principle we have

$$u(t + \tau_n', x) \geqslant W_{\delta_2}(x - (\beta - c_0 + \delta_2)t - X') \text{ for } x \in J_2(t),\ t > 0,$$

where $J_2(t) := [(\beta - c_0 + \delta_2)t + X' - L_{\delta_2}, (\beta - c_0 + \delta_2)t + X']$. By $\psi_n^{(2)}(\tau_n') = 0$ we have

$$y_n = (\beta - c_0 + \delta_2)(t_n - \tau_n') + X' - z_{\delta_2} \in J_2(t_n - \tau_n').$$

Hence at $t = t_n - \tau_n'$ and $x = y_n$ we have

$$1 - \varepsilon > u(t_n, y_n) \geqslant W_{\delta_2}(y_n - (\beta - c_0 + \delta_2)(t_n - \tau_n') - X') = W_{\delta_2}(-z_{\delta_2}) = D_{\delta_2} > 1 - \varepsilon,$$

a contradiction.

This completes the proof of Theorem 2.5. □

References

[1] S.B. Angenent, The zero set of a solution of a parabolic equation, J. Reine Angew. Math. 390 (1988) 79–96.

[2] D.G. Aronson, H.F. Weinberger, Multidimensional nonlinear diffusions arising in population genetics, Adv. Math. 30 (1978) 33–76.

[3] I.E. Averill, The effect of intermediate advection on two competing species, Doctor of Philosophy, Ohio State University, Mathematics, 2012.

[4] M. Ballyk, H. Smith, A model of microbial growth in a plug flow reactor with wall attachment, Math. Biosci. 158 (1999) 95–126.

[5] F. Belgacem, C. Cosner, The effect of dispersal along environmental gradients on the dynamics of populations in heterogeneous environment, Can. Appl. Math. Q. 3 (1995) 379–397.

[6] H. Berestycki, F. Hamel, Front propagation in periodic excitable media, Comm. Pure Appl. Math. 55 (8) (2002) 949–1032.

[7] G. Bunting, Y. Du, K. Krakowski, Spreading speed revisited: analysis of a free boundary model, Netw. Heterog. Media 7 (4) (2012) 583–603.

[8] J. Byers, J. Pringle, Going against the flow: retention, range limits and invasions in advective environments, Mar. Ecol. Prog. Ser. 313 (2006) 27–41.

[9] X. Chen, A. Friedman, A free boundary problem arising in a model of wound healing, SIAM J. Math. Anal. 32 (2001) 778–800.

[10] Y. Du, Z.M. Guo, Spreading–vanishing dichotomy in the diffusive logistic model with a free boundary II, J. Differential Equations 250 (2011) 4336–4366.
[11] Y. Du, Z.M. Guo, The Stefan problem for the Fisher-KPP equation, J. Differential Equations 253 (2012) 996–1035.
[12] Y. Du, Z. Lin, Spreading–vanishing dichotomy in the diffusive logistic model with a free boundary, SIAM J. Math. Anal. 42 (2010) 377–405.
[13] Y. Du, B. Lou, Spreading and vanishing in nonlinear diffusion problems with free boundaries, J. Eur. Math. Soc. (JEMS) (2015), in press, arXiv:1301.5373.
[14] Y. Du, B. Lou, M. Zhou, Nonlinear diffusion problems with free boundaries: convergence, transition speed and zero number arguments, SIAM J. Math. Anal. (2015), to appear.
[15] Y. Du, H. Matano, K. Wang, Regularity and asymptotic behavior of nonlinear Stefan problems, Arch. Ration. Mech. Anal. 212 (2014) 957–1010.
[16] Y. Du, H. Matsuzawa, M. Zhou, Sharp estimate of the spreading speed determined by nonlinear free boundary problems, SIAM J. Math. Anal. 46 (2014) 375–396.
[17] A. Ducrot, T. Giletti, H. Matano, Existence and convergence to a propagating terrace in one-dimensional reaction–diffusion equations, Trans. Amer. Math. Soc. 366 (2014) 5541–5566.
[18] F.J. Fernandez, Unique continuation for parabolic operators. II, Comm. Partial Differential Equations 28 (2003) 1597–1604.
[19] H. Gu, Z. Lin, B. Lou, Long time behavior of solutions of a diffusion–advection logistic model with free boundaries, Appl. Math. Lett. 37 (2014) 49–53.
[20] H. Gu, Z. Lin, B. Lou, Different asymptotic spreading speeds induced by advection in a diffusion problem with free boundaries, Proc. Amer. Math. Soc. 143 (2015) 1109–1117.
[21] F. Hamel, J. Nolen, J. Roquejoffre, L. Ryzhik, A short proof of the logarithmic Bramson correction in Fisher-KPP equations, Netw. Heterog. Media 8 (2013) 275–289.
[22] D. Hilhorst, M. Iida, M. Mimura, H. Ninomiya, A competition–diffusion system approximation to the classical two-phase Stefan problem, Jpn. J. Ind. Appl. Math. 18 (2001) 161–180.
[23] D. Hilhorst, M. Mimura, R. Schätzle, Vanishing latent heat limit in a Stefan-like problem arising in biology, Nonlinear Anal. Real World Appl. 4 (2003) 261–285.
[24] S.B. Hsu, Y. Lou, Single phytoplankton species growth with light and advection in a water column, SIAM J. Appl. Math. 70 (2010) 2942–2974.
[25] J. Huisman, M. Arrayas, U. Ebert, B. Sommeijer, How do sinking phytoplankton species manage to persist?, Amer. Nat. 159 (2002) 245–254.
[26] Y. Kanako, H. Matsuzawa, Spreading speed and sharp asymptotic profiles of solutions in free boundary problems for nonlinear advection–diffusion equations, J. Math. Anal. Appl. 428 (2015) 43–76.
[27] Y. Kaneko, Y. Yamada, A free boundary problem for a reaction–diffusion equation appearing in ecology, Adv. Math. Sci. Appl. 21 (2011) 467–492.
[28] X. Liu, B. Lou, Asymptotic behavior of solutions to diffusion problems with Robin and free boundary conditions, Math. Model. Nat. Phenom. 8 (2013) 18–32.
[29] X. Liu, B. Lou, Long time behavior of solutions of a reaction–diffusion equation with Robin and free boundary conditions, J. Differential Equations 259 (2015) 423–453.
[30] N.A. Maidana, H. Yang, Spatial spreading of West Nile Virus described by traveling waves, J. Theoret. Biol. 258 (2009) 403–417.
[31] I.G. Petrovski, Ordinary Differential Equations, Prentice Hall, Englewood Cliffs, NJ, 1966; Dover, New York, 1973.
[32] A. Potapov, M. Lewis, Climate and competition: the effect of moving range boundaries on habitat invasibility, Bull. Math. Biol. 66 (5) (2004) 975–1008.
[33] G.A. Riley, H. Stommel, D.F. Bumpus, Quantitative ecology of the plankton of the western North Atlantic, Bull. Bingham Oceanogr. Collect. Yale Univ. 12 (1949) 1–169.
[34] L.I. Rubinstein, The Stefan Problem, American Mathematical Society, Providence, RI, 1971.
[35] N. Shigesada, A. Okubo, Analysis of the self-shading effect on algal vertical distribution in natural waters, J. Math. Biol. 12 (1981) 311–326.
[36] D.C. Speirs, W.S.C. Gurney, Population persistence in rivers and estuaries, Ecology 8 (5) (2001) 1219–1237.
[37] O. Vasilyeva, F. Lutscher, Population dynamics in rivers: analysis of steady states, Can. Appl. Math. Q. 18 (4) (2010) 439–469.
[38] M.X. Wang, The diffusive logistic equation with a free boundary and sign-changing coefficient, J. Differential Equations 258 (2015) 1252–1266.
[39] P. Zhou, D.M. Xiao, The diffusive logistic model with a free boundary in heterogeneous environment, J. Differential Equations 256 (2014) 1927–1954.